

Pre-conception serum ferritin concentrations are associated with metal concentrations in blood during pregnancy: A cohort study in Benin

Sarah Davies, Valérie Briand, Manfred Accrombess, Nadine Fievet, Barbara Le Bot, Séverine Durand, Ginoa Gbota, Emmanuel Yovo, Bertin Vianou,

Darius Sossou, et al.

▶ To cite this version:

Sarah Davies, Valérie Briand, Manfred Accrombess, Nadine Fievet, Barbara Le Bot, et al.. Preconception serum ferritin concentrations are associated with metal concentrations in blood during pregnancy: A cohort study in Benin. Environmental Research, 2021, 202, 10.1016/j.envres.2021.111629. hal-03312406

HAL Id: hal-03312406 https://ehesp.hal.science/hal-03312406

Submitted on 2 Aug 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

Version of Record: https://www.sciencedirect.com/science/article/pii/S0013935121009233 Manuscript_da92bf56e237689c3cb916fbba55cd4f

1 Pre-conception serum ferritin concentrations are associated with metal concentrations in blood

2 during pregnancy: A cohort study in Benin

- 3 Sarah Davies^{a,b} (sdavies72@gmail.com), Valérie Briand^c (valerie.briand@ird.fr), Manfred
- 4 Accrombessi^{d,e} (accrombessimanfred@yahoo.fr), Nadine Fievet^{d,f} (nadine.fievet@ird.fr), Barbara Le
- 5 Bot^g (Barbara.LeBot@ehesp.fr), Séverine Durand^g (severine.durand@ehesp.fr), Gino Agbota^d
- 6 (ginodric@yahoo.fr), Emmanuel Yovo^d (emmanuelkoffiyovo@yahoo.fr), Bertin Vianou^d
- 7 (bertinosidase@yahoo.fr), Darius Sossou^d (anax20032002@yahoo.fr), Yves Martin-Prevelⁱ
- 8 (yves.martin-prevel@ird.fr), Achille Massougbodji^h (massougbodjiachille@yahoo.fr), Michel Cot^f
- 9 (michel.cot@ird.fr), Philippe Glorennec^{*g*} (philippe.glorennec@ehesp.fr), Florence Bodeau-Livinec^{*a,b*}
- 10 (florence.bodeau-livinec@ehesp.fr)
- 11 ^a Département Méthodes Quantitatives en Santé Publique (METIS), Ecole des Hautes Etudes en Santé
- 12 Publique (EHESP), F-35000 Rennes, France
- 13 ^b Université de Paris, Center of Research in Epidemiology and Statistics/CRESS, INSERM, INRA,
- Obstetrical, Perinatal and Pediatric Epidemiology Research Team (EPOPé)—UMR1153, F-75004,
 Paris, France
- ^c Institut de Recherche Pour le Développement (IRD), University of Bordeaux, Inserm, UMR 1219, 146
- 17 rue Léo-Saignat, 33076, Bordeaux Cedex, France, Université de Paris, MERIT, IRD, 75006, Paris,
- 18 France
- ^d Institut de Recherche Clinique du Benin (IRCB), 04 BP1114, Abomey-Calavi, Benin
- 20 ^e Faculty of Infectious and Tropical Diseases, Disease Control Department, London School of Hygiene
- 21 and Tropical Medicine, London, United Kingdom
- ^fInstitut de Recherche pour le Développement (IRD), Paris Descartes Université, 75006 Paris, France
- ⁹ Ecole des Hautes Études en Santé Publique (EHESP), Institut National de la Sante et de la Recherche
- 24 Medicale (Inserm), Institut de Recherche en Santé, Environnement et Travail (Irset)—UMR_S 1085,
- 25 University of Rennes, F-35000 Rennes, France
- 26 ^h University of Abomey-Calavi Faculté des Sciences de la Santé, Cotonou, Bénin
- 27 ^{*i*} Nutripass Research Unit, Institut de Recherche pour le Développement (IRD), University of
- 28 Montpellier, SupAgro, Montpellier, France

29 Corresponding Author Contact Information

- 30 Name: Sarah Davies
- 31 E-mail : sdavies72@gmail.com
- 32 Present Postal address : 24 Gow St, Hamilton North 2292, Australia
- 33 Sarah Davies: Methodology, Formal analysis, Writing Original Draft Valérie Briand:
- 34 Conceptualization, Supervision, Writing Review & Editing Manfred Accrombessi: Investigation,
- 35 Writing Review & Editing Nadine Fievet: Conceptualization, Supervision Barbara Le Bot:
- 36 Investigation, Formal Analysis Séverine Durand: Investigation, Formal Analysis, Writing Review &
- 37 Editing **Gino Agbota:** Investigation, Writing Review & Editing **Emmanuel Yovo:** Investigation **Bertin**
- 38 Vianou: Biological Analysis Darius Sossou: Biological Analysis Yves Martin-Prevel: Conceptualization,
- 39 Supervision, Writing Review & Editing Achille Massougbodjii: Supervision Michel Cot: Supervision
- 40 **Philippe Glorennec:** Methodology, Writing Review & Editing **Florence Bodeau-Livinec:**
- 41 Conceptualization, Methodology, Supervision, Writing Review & Editing

42 Title

43 Pre-conception serum ferritin concentrations are associated with metal concentrations in 44 blood during pregnancy: A cohort study in Benin

45 Abstract

- 46 **Background:** Iron deficiency is a common nutritional deficiency that impacts maternal health and
- 47 fetal development and is also associated with increased uptake of toxic metals. Women in sub-
- 48 Saharan Africa are highly exposed to both iron deficiency and metals in the environment. As research
- 49 on the developmental origins of health and disease increasingly shows impacts of pre-conception
- 50 maternal health on pregnancy and fetal health, these environmental exposures are of concern.
- 51 **Objectives:** This study investigated the association between iron status pre-pregnancy and blood
- 52 metal concentrations in the first trimester of pregnancy with potential implications for iron
- 53 supplementation.
- 54 **Methods**: Pre-conception and first trimester blood samples taken from 262 Beninese women were
- 55 tested for serum ferritin, inflammation markers, manganese (Mn), cadmium (Cd), lead (Pb), copper,
- 56 zinc, selenium, mercury and arsenic. Associations between serum ferritin adjusted for inflammation
- 57 and metal concentrations were analyzed using multivariate linear regression.
- 58 **Results:** Women with iron deficiency before conception (13%) were more likely to remain iron
- deficient in the first trimester (4%) (adjusted OR=41.2, 95%CI 6.2; 275.0) even within the context of
 routine iron supplementation during pregnancy.
- 61 Lower pre-pregnancy serum ferritin concentrations were significantly related to higher
- 62 concentrations of Mn, Cd and Pb in the first trimester. Every 1% increase in serum ferritin
- 63 concentration was associated with a 0.13% decrease in Mn (adjusted β =-0.13, 95%Cl -0.18; -0.07), a
- 64 0.22% decrease in Cd (adjusted β =-0.22, 95%Cl -0.28; -0.15) and a 0.06% decrease in Pb
- 65 concentration (adjusted β =-0.06, 95%Cl -0.12; -0.006).
- 66 **Discussion:** These results suggest that increasing iron stores prior to pregnancy may prevent
- 67 excessive uptake of toxic concentrations of the metals Mn, Cd and Pb and argue in favour of testing
- 68 the effects of iron supplementation prior to pregnancy on metal concentrations.
- 69 **Keywords:** Iron deficiency, metals, environmental exposure, pregnancy, pre-conception

70

71

- 72 **Funding Acknowledgement:** This study was supported by the French National Research Agency
- 73 (ANR, ANR-13-JSV1-0004, grant 2013), the Fondation de France (no: 00074147, grant 2017) and a
- 74 grant from the Ecole des Hautes Etudes en Santé Publique (ESSOR)
- 75 **Competing Interests:** The authors declare they have no actual or potential competing financial
- 76 interests or personal relationships that could have appeared to influence the work reported in this 77 paper.
- 78 **Ethics approval**: The authors confirm that consent from subjects participating in the cohort study
- 79 was received prior to conducting the study and that the study was approved by the Ethics Committee
- 80 of the Institut des Sciences Biomédicales Appliquées (ISBA) in Benin.

81

82 1. Introduction

- 83 Iron deficiency is the most common nutritional deficiency worldwide and with or without anemia has
- 84 important consequences for human health and child development. Globally, about 33% of all women
- of reproductive age are estimated to be anemic (WHO, 2016a), with 16-25% estimated to be due to
- 86 iron deficiency (Petry et al., 2016). Women's iron requirements increase significantly during
- 87 pregnancy, rising during the second trimester and continuing to increase throughout the remainder
- 88 of pregnancy (Bothwell, 2000).
- 89 Iron is essential for brain development, particularly in late fetal and early postnatal life, where
- 90 deficits during these critical periods can permanently affect brain function (Georgieff et al., 2018).
- 91 Maternal iron deficiency also reduces iron stores in the fetus which further exposes neonates to
- 92 neurodevelopmental risks as infants cannot regulate iron absorption until 6-9 months of age
- 93 (Radlowski & Johnson, 2013). The WHO recommends oral iron and folic acid supplementation for
- 94 pregnant women to prevent maternal anaemia, puerperal sepsis, low birth weight, and preterm birth
- 95 (WHO, 2016b).
- 96 Higher iron levels, however may be detrimental in malaria endemic areas such as Benin, as some
- 97 studies in children have found malaria risk increases in relation to serum ferritin concentrations and
- 98 iron deficiency seems to act as a protective factor (Moya-Alvarez et al., 2017; Muriuki et al., 2019). A
- 99 recent systematic review of malaria risk and iron status in pregnancy found that iron deficiency was
- 100 only associated with increased risk of malaria when measured by serum ferritin and not when other
- biomarkers of iron deficiency were included. In the light of limited evidence however, iron
- 102 supplementation is still recommended to be provided in combination with malaria prevention during
- 103 pregnancy (Sangaré et al., 2014).
- 104 Iron deficiency is also associated with the increased uptake of divalent metals which can adversely
 affect neurodevelopment (Kim, 2018). Dietary non-heme iron is transported into intestinal cells by
 the divalent metal-ion transporter 1 (DMT1) and then exported into circulation via ferroportin 1
- 107 (FPN1) or stored as ferritin. Iron homeostasis is regulated by the hormone hepcidin, which is in low
- 108 concentrations when tissue iron stores are low, up-regulating DMT1 and FPN1 (Anderson & Frazer,
- 2017). DMT1 has been demonstrated to transport divalent metal ions other than iron, notably
- 110 manganese, cadmium and lead as well as to a lesser extent copper and zinc (Bressler et al.,2004;
- 111 Garrick et al., 2006). There is also evidence that FPN1 can transport manganese and zinc (Loréal et
- 112 al., 2014; Madejczyk & Ballatori, 2012).
- 113 Research conducted within the framework of the Developmental origins of Health and Disease
- 114 (DoHAD) continues to provide evidence of the long-term consequences on health of environmental
- exposures from conception and including the role of maternal nutrition prior to conception (Mandy
- 8 Nyirenda, 2018; Stephenson et al., 2018). Many studies have examined the effects of iron
- deficiency and its relationship to a range of environmental metal exposures during pregnancy and in
- relation to fetal and childhood development. This study examines data from a unique cohort where
- 119 women were recruited prior to conception, allowing iron status before pregnancy to be examined in
- 120 relation to whole blood concentrations in early pregnancy of 8 potentially toxic metals: Manganese,
- 121 Copper, Zinc, Selenium, Lead, Cadmium, Mercury and Arsenic.

- 122 Manganese is an essential nutrient but also a neurotoxicant at high exposure levels. Lead and
- 123 cadmium are long-lived, toxic heavy metals that have no known beneficial function in the human
- 124 body. The presence of these metals in the environment and the likelihood of increased uptake before
- 125 and during pregnancy for women with low iron stores can result in a range of deleterious effects.
- 126 Cd accumulation over time can damage renal function and affect bone metabolism (ATSDR, 2012a)
- as well as some evidence it is toxic to endocrine systems resulting in decreased birth weights
- 128 (Rahman et al., 2016). There is some evidence that prenatal Cd exposure may impact cognitive
- development in children (Kippler et al., 2012). Although Cd accumulates in the placenta acting as a
- 130 partial barrier to foetal exposure with cord blood concentrations significantly lower than maternal
- blood (Esteban-Vasallo María D. et al., 2012; Truska et al., 1989), Cd also affects placental transport
- 132 of calcium and zinc which are essential for foetal development (Gundacker & Hengstschläger, 2012).
- 133 The health effects of Pb are diverse, not only is Pb a known developmental neurotoxicant but
- exposure to Pb is associated with toxicity to every organ system (ATSDR, 2020). Lead readily crosses
- the placenta (Bellinger, 2005; Rossipal et al., 2000) and there is good evidence that maternal lead
- 136 exposure during pregnancy can adversely affect both maternal and child health. There is no apparent
- 137 threshold below which adverse effects of lead do not occur (Budtz-Jørgensen et al., 2013; CDC,
- 138 2010). Bone lead stores persist for decades and are mobilized during periods of increased bone
- 139 turnover such as pregnancy and lactation. Iron status, along with intake of dietary calcium and zinc
- 140 can affect absorption of Pb, potentially leading to alterations in blood lead concentrations (ATSDR,
- 141 2020). In addition, there is some evidence that combined Pb and Mn exposures in-utero and in
- 142 childhood could aggravate negative impacts on neurodevelopment (Kim et al., 2009; Lin et al., 2013;
- 143 Menezes-Filho et al., 2018).
- Se is an essential trace element which can also be toxic and has a complex interplay of metabolic
- 145 interactions with numerous other nutrients and toxic substances (Arnaud & van Dael, 2018; Sun et
- al., 2014). Se intake varies widely dependent on concentrations in soils transferred to crops.
- 147 The objective of this study was to investigate the association between iron status pre-pregnancy and
- 148 during the first trimester with the blood concentrations of eight metals in the first trimester of
- 149 pregnancy: manganese, cadmium, lead, copper, zinc, selenium, mercury and arsenic.
- 150

151 2. Methods

152 2.1 Study Population

This study includes 262 women from the REtard de Croissance Intra-uterin et PALudisme (RECIPAL) cohort in southern Benin primarily designed to assess the consequences of malaria in early pregnancy for the fetus as detailed in Accrombessi et al., 2018 and summarized in Figure 1. Women in the RECIPAL cohort were recruited prior to conception between June 2014 and December 2016 and followed monthly until pregnancy and through to delivery. At inclusion demographic, household and socioeconomic characteristics were collected. In addition, blood samples were taken at

- 159 enrolment before conception and at the second antenatal care (ANC2) visit which occurred for most
- 160 women during the 1st trimester of pregnancy.

161 2.2 Ethics Statement

- 162 The RECIPAL study was approved by the Ethics Committee of the Institut des Sciences Biomédicales
- 163 Appliquées (ISBA) in Benin. Women enrolled in the cohort received regular free medical follow-up
- 164 treatment as necessary during the life of the study. Blood samples were stored and assessed for
- 165 metals at a later date, these results at the population level were presented to health authorities at
- 166 local and national levels.
- 167

168 **Figure 1:** Flowchart diagram of the study



169

170 2.3 Gestational Follow-up

- 171 According to Beninese national recommendations, a kit including iron and folic acid tablets for 1
- 172 month was given to each woman at the first ANC visit and maternity staff were encouraged to
- administer iron and folic acid supplementation throughout the pregnancy (200 mg oral ferrous
- sulfate and 10 mg folic acid daily). As part of the study, the cost of drugs were supported by the
- project; however, the iron and folic acid supplementation were not observed by the study staff.

176 2.4 Blood Measurements

177 Blood samples were initially tested for parameters related to iron deficiency, inflammation markers 178 and malaria. Blood serum samples were tested for ferritin, C-reactive protein (CRP) and α_1 -acid 179 glycoprotein (AGP). Ferritin may be measured in either serum or plasma, with both positively 180 correlated with the size of total body iron stores (WHO, 2020). As nutrition was one of the core 181 issues in the RECIPAL cohort, serum ferritin was used as part of the assessment of nutritional status 182 and the micronutrient intake of women. Hb level was measured with a HemoCue. Before conception and during pregnancy, malarial screening was performed using both microscopy (thick blood smear) 183 184 and PCR. Subsequently two sets of ANC2 whole blood samples of women who completed the study 185 follow-up were tested for metals. A first group of 60 tested for Manganese (Mn), Copper (Cu), Zinc 186 (Zn), lead (Pb) and Cadmium (Cd) with results reported previously (Guy et al., 2018). Samples from a 187 further 202 were later tested for these five metals with the addition of tests for Selenium (Se), 188 Mercury (Hg) and Arsenic (As).

189 The 262 whole blood samples stored at -18°C, were analyzed by a plasma torch coupled with tandem 190 mass spectrometer (ICP-MSMS 8800, Agilent Technologie) after mineralization. Mineralization is 191 performed by adding nitric acid and heating at 83°C for 4 hours using a graphite block (Hotblock Pro, 192 model SC-189, Environmental Express). The mineralization diluted at 1:20 was then analyzed, the use 193 of the tandem mass spectrometer and the oxygen mode made it possible to manage the 194 interferences in particular for the determination of selenium and arsenic. An online addition of 195 internal standards (Sc, Ge, 77Se, Rh, Re and Ir) was performed to correct matrix effects and to verify 196 the absence of drift (signal variation less than 25%). The quantification of Se was performed by 197 isotope dilution with 77Se to resolve persistent effects. At each analysis sequence, consumable 198 blanks and certified blood controls were systematically analyzed (Utak level 1, Utak level 2). The 199 results of the blood tests were validated if the concentrations of the blood controls were within the 200 limits of the control cards. The method of analysis was validated by 2 good inter-laboratory tests over 201 1 year. This method is accredited by the French accreditation committee (Cofrac). The limits of 202 quantification (LOQ) of Pb, Mn, Cu, Zn, Cd, As, Se and Hg are respectively 2, 5, 200, 200, 0.25, 1, 20 203 and 0.4 μ g/L.

204 2.5 Definitions

205 *Iron deficiency* was defined as serum ferritin less than 15 μ g/l after adjustment for inflammation. 206 Serum ferritin is a reliable indicator of total body iron stores when inflammation is not present. In 207 regions such as the study area in Benin where there is widespread infection or inflammation, defining 208 iron deficiency using serum ferritin becomes complicated as ferritin is a positive acute phase 209 response protein. WHO recommends in these areas to assess serum ferritin with concurrent 210 measurement of CRP and AGP and apply one of four possible methods: raising the cut-off value that 211 defines iron deficiency to 70 µg/l; excluding individuals with elevated CRP or AGP concentrations; use 212 an arithmetic correction; or use a regression correction (WHO, 2020). A comparison of methods on a 213 cohort of Kenyan pregnant women concluded that using a regression correction approach led to 214 better estimates of iron deficiency (Mwangi et al., 2019). This regression correction approach 215 according to the following equation was also used in this study as it allowed a correction to

- continuous data and the inclusion of a correction for inflammation present due to malaria infection.
- 217 (Namaste et al., 2017; WHO, 2020).
- 218 Ferritin_{adjusted} = Ferritin_{unadjusted} $\beta_1(CRP_{obs} CRP_{ref}) \beta_2(AGP_{obs} AGP_{ref}) \beta_3 malaria$
- where β_x are the regression coefficients, obs is the observed value and ref is the external reference value generated to define low inflammation.
- 221 Inflammation was defined as either CRP > 5 mg/l and/or AGP > 1 g/l (Namaste et al., 2017; WHO,
 222 2020)
- 223 Anemia was defined prior to pregnancy as hemoglobin <120 g/l and during pregnancy as <110 g/l
- with *moderate anemia* between 70 to 110 g/l and *severe anemia* < 70 g/l (WHO, 2015)
- 225 Iron deficiency anemia was defined as those meeting the criteria both for iron deficiency and anemia.
- 226 *Insufficient iron stores for pregnancy* was defined as serum ferritin <32 μg/l approximately equivalent
- to 300 mg of total body iron stores, the estimated minimum with which a woman must enter
- 228 pregnancy to meet her requirements fully assuming an optimal diet.
- 229 First trimester was defined as less than 14 weeks gestation based on both ultrasound scan and LMP

230 2.6 Statistical Analysis

231 The association between serum ferritin pre-pregnancy and at ANC2 and blood metal concentrations 232 for Mn, Cu, Zn, Pb and Cd, Se, Hg and As were modelled separately using multivariate linear 233 regression. Log transformations of the metal concentrations were made due to non-normal 234 distributions except for Se which did not have a distribution significantly different from normal when 235 using a Shapiro-Wilk test. Log transformations of serum ferritin concentrations were also used in the 236 models for all metals except Se as this improved the linear relationships checked by visual inspection. 237 Confounders were selected based on literature review, the consideration of directed acyclic graphs 238 and after testing for collinearity between candidate variables. Gestational age at ANC2, gravidity and 239 indicators for education level and wealth were included in the models for all metals. As there was 240 little variability in education level, literacy was used as the variable to represent whether any formal 241 education had been received. At inclusion in the cohort, detailed questions were asked on the 242 ownership of assets, the number owned was used as an indicator of wealth. Other potential 243 confounders were tested such as age, body mass index (BMI), the time between pre-pregnancy and 244 ANC2 testing, being positive for malaria by microscopy at ANC2 and living lakeside or inland as a 245 potential indicator of different environmental and dietary exposures. Univariate analysis was first 246 conducted retaining variables with a p value less than 0.25. These variables were then removed 247 stepwise from multivariate models if p values were greater than 0.05. Analysis was carried out using 248 Stata version 15 for Windows.

249 3. Results

250 3.1 Population characteristics

251 Table 1 presents the general characteristics of the 262 women included in the study population. 252 Women were on average 27 years old, primarily of Toffin ethinicity (74.4%) and living within 4 sub-253 districts of Sô-Ava district: Vekky and Houedo-Aguekon on the edge of the polluted Lake Nokoue; Sô-254 Ava along the River So; and Akassato where women lived within 3 to 10 km from Lake Nokoue. The 255 majority of women were multigravidae (92.4%), were non-smokers (99.2%) and were within a 256 healthy BMI range at the start of pregnancy (67%). Table 2 presents the blood test results related to 257 iron deficiency at pre-pregnancy and ANC2 visits and Table 3 gestational age information at the first 258 two ANC visits. Before pregnancy, 12.7% of women were iron deficient, 57.9% anemic and 7% tested 259 positive for malaria. Most women became pregnant within 6 months of inclusion (51.2%) with a 260 further 37.3% conceiving within a year. Pregnancy was confirmed at the first ANC visit which 261 occurred at an average gestational age of 6.9 weeks at which time iron supplementation was 262 prescribed. Gestational age at ANC2, at which blood was sampled for iron parameters and metals 263 analysis, averaged 11.7 weeks with 84.4% considered in the first trimester. By ANC2 only 3.5% of 264 women were still considered as iron deficient, 38.9% anemic and 5.4% tested positive for malaria.

265 **Table 1:** Study Population Descriptive Statistics

Maternal Characteristics	No.	Mean ± SD, %
Age	262	26.7 (±4.9)
Subdistrict		
Vekky	92	35.1%
Sô-Ava	91	34.7%
Akassato	65	24.8%
Houedo-Aguekon	14	5.3%
Marital Status		
Married (Monogamy)	173	66.0%
Married (Polygamy)	71	27.1%
Cohabitation	18	6.9%
Ethnicity		
Toffin	195	74.4%
Aizo	32	12.2%
Fon	20	7.6%
Other	13	5.8%
Household density ^a	262	6.1 (±3.9)
Education		
Illiterate	183	69.8%
Gravidity		
Primigravidae	20	7.6%
Multigravidae	242	92.4%
Tobacco during pregnancy		
During 1st trimester	1	0.4%
Throughout pregnancy	1	0.4%
None	254	99.2%
BMI in early pregnancy	l	
Underweight (<18.5)	24	9.2%
Normal (18.5-25)	175	66.8%

	-			
	Overweight/Obese (≥25)	63	24.0%	
266	^a Number of people per household			

267 **Table 2:** Maternal blood test results for iron deficiency parameters

	No.	Pre-pregnancy visit	No.	2nd Ante Natal Care
Hemoglobin (g/l)				(ANCZ) VISIL
	264	447 (1420)	200	
Mean (±SD)	261	117.6 (±13.0)	260	114.1 (±11.8)
Range (min - max)		84 - 151		84 - 146
Ferritin (µg/l)				
Mean (±SD)	260	58.1 (±34.6)	261	84.0 (±40.7)
Range		4.3 - 208		10.3 - 187
Adjusted Ferritin (µg/l) ^a				
Mean (±SD)	260	42.0 (±23.6)	261	74.1 (±35.8)
Range		4.3 - 105		9.9 - 171
Anemia ^b	261	151 (57.9%)	260	101 (38.9%)
Iron Deficiency ^c	260	33 (12.7%)	261	9 (3.5%)
Iron Deficiency Anemia ^d	260	25 (9.6%)	261	5 (1.9%)
Inflammation present ^e	260	48 (18.5%)	261	63 (24.1%)
Malaria positive by microscopy	259	18 (7.0%)	261	14 (5.4%)
Insufficient iron stores for pregnancy ^f	260	98 (37.7%)		

268 ^{*a*}Ferritin results adjusted for inflammation based on C-Reactive Protein (CRP) and α_1 -acid-glycoprotein (AGP) biomarkers

according to the BRINDA approach

270 *b*Anemia defined pre-pregnancy as hemoglobin <120g/l and during pregnancy as <110g/l

 $271 \qquad {}^{c} Iron \ deficiency \ defined \ as \ adjusted \ serum \ ferritin \ <15 \ \mu g/L$

272 dIron deficiency anemia defined as meeting both anemia and iron deficiency criteria

- $273 \qquad {}^{e} Inflammation \ defined \ as \ either \ CRP>5 \ \mu g/l \ or \ AGP>1 \ \mu g/l$
- 274 $f Insufficient iron stores for pregnancy defined as adjusted serum ferritin <32 $\mu g/L$

275

276 **Table 3:** Gestational age information and maternal blood test timing

	No.	1st Ante Natal Care (ANC1) visit ^g	No.	2nd Ante Natal Care (ANC2) visit
Gestational age (weeks)				
Mean (±SD)	262	6.9 (±2.3)	261	11.7 (±2.8)
Range		3.7 – 19.3		7.0 – 24.0
1 st Trimester (≤14 weeks gestation)			262	221 (84.4%)
2 nd Trimester (>14 weeks gestation)			262	41 (15.6%)
Time from pre-pregnancy blood test to confirmation of pregnancy (weeks)	260			
Geometric Mean (95%CI)		17.4 (15.6; 19.4)		
Range		2.1 - 106		
n (%) > 26 weeks		97 (37.3%)		
n (%) > 52 weeks		30 (11.5%)		

277

⁹Pregnancy confirmed at the first ante-natal visit and iron supplements provided to the mother from this point onwards

278

279 3.2 Metal Associations

- 280 The range of concentrations in maternal blood at the ANC2 visit for the eight metals tested is shown
- 281 in Figure 2. The median and interquartile range concentrations were for Mn 11.2 μg/l (IQR 8.76-
- 282 13.2), Cu 1333 μg/l (IQR 1165-1490), Zn 5074 μg/l (IQR 4541-5830), Se 149.1 μg/l (IQR 132.1-167.3),
- 283 Pb 31.3 μ g/l (IQR 25.7-38.9), Cd 0.253 μ g/l (IQR 0.177-0.331), and Hg 1.20 μ g/l (IQR 0.907-1.56).

- 284 Figure 2: Maternal blood concentrations of elements (μg/l) at ANC2 for n=262 for Mn, Cu, Zn, Pb, Cd
- 285 and n=202 for Se, Hg, As^{*}



*Boxes represent median values with interquartile range (IQR) and whiskers highest and lowest points within 1.5 IQR length
 from the box. Dots represent outliers

288 The relationship between pre-pregnancy iron deficiency and iron deficiency at ANC2 is presented in

Table 4. Women found to be iron deficient before conception were significantly more likely to remain

iron deficient when re-tested in the first trimester (adjusted OR=41.2, 95%Cl 6.18; 275). However,

the numbers of iron deficient women, were small with 13% of the sample found to be iron deficient

before conception and 4% in the first trimester.

293 **Table 4:** Effect of pre-pregnancy iron deficiency on iron deficiency at ANC2

	n	Crude OR (95% CI)	Adjusted OR (95% CI) ^a
		Iron defic	ient ^b at ANC2
Iron deficient ^b pre-pregnancy	259	16.5 (3.24;106)	41.2 (6.18;275)
		Iron stores insufficient ^c at ANC2	
Iron stores insufficient for pregnancy ^c at pre-pregnancy blood test	259	10.3 (4.23; 27.2)	12.5 (5.11;30.6)

- ^aAdjusted for G2 gestational age, gravidity, education and wealth (based on no. possessions)
- ^bIron deficiency defined as serum ferritin adjusted for inflammation <15 μg/L
- 296 cInsufficient iron stores for pregnancy based on an estimated requirement of at least 300mg which corresponds to a serum 297 ferritin level of \geq 32 µg/L. As iron requirements are low in the first trimester, the same cut-off was used for pre-pregnancy
- and ANC2 blood test results
- 299
- 300 The results for blood metal concentrations at ANC2 showed that higher pre-pregnancy serum ferritin 301 concentrations and higher ANC2 serum ferritin concentrations were significantly related to lower 302 concentrations of Mn, Cd and Pb in the first trimester after adjustment for potential confounders 303 (Table 5). Every 1% increase in pre-pregnancy serum ferritin concentration was associated with a 304 0.13% decrease in Mn (adjusted β =-0.13, 95%Cl -0.18; -0.07), a 0.22% decrease in Cd (adjusted β =-305 0.22, 95%CI -0.28; -0.15) and a 0.06% decrease in Pb concentration (adjusted β =-0.06, 95%CI -0.12; -306 0.006). Similarly, every 1% increase in ANC2 serum ferritin concentration was associated with a 307 0.19% decrease in Mn (adjusted β =-0.19, 95%Cl -0.25; -0.13), a 0.21% decrease in Cd (adjusted β =-
- 308 0.21, 95%CI -0.29; -0.13) and a 0.11% decrease in Pb concentration (adjusted β =-0.11, 95%CI -0.17; -
- 309 0.04).
- 310 The only other metal to show a significant positive association was Se where every 1 microgram unit
- 311 increase in pre-pregnancy serum ferritin concentration was associated with a 0.19 microgram
- increase in Se concentration (adjusted β =0.19, 95%CI 0.043; 0.34) and every 1 unit microgram
- increase in ANC2 serum ferritin concentration was associated with a 0.11 microgram increase in Se
- 314 concentration (adjusted β =0.11, 95%CI 0.013; 0.22).
- 315 After adjusting the models no significant associations were found between serum ferritin
- 316 concentrations and Zn, Cu, As or Hg blood concentrations in the first trimester.
- **Table 5:** Associations between serum ferritin concentrations pre-pregnancy and ANC2 blood metal
- 318 concentrations

	No.	Crude β (95% Cl)	Adjusted β (95% Cl)
Log Mn (μg/l)			
Log Serum ferritin pre-pregnancy (µg/l)	259	-0.13 (-0.18, -0.07)	-0.13 (-0.18, -0.07) ^a
Log Serum ferritin at ANC2 (μg/l)	259	-0.19 (-0.25, -0.13)	-0.19 (-0.25, -0.13) ^a
Log Cd (µg/l)			
Log Serum ferritin pre-pregnancy (µg/l)	259	-0.21 (-0.28, -0.15)	-0.22 (-0.28, -0.15) ^b
Log Serum ferritin at ANC2 (μg/l)	260	-0.20 (-0.28, -0.13)	-0.21 (-0.29, -0.13) ^b
Log Zn (μg/l)			
Log Serum ferritin pre-pregnancy (µg/l)	259	-0.03 (-0.07, 0.01)	-0.03 (-0.07, 0.01) ^b
Log Serum ferritin at ANC2 (μg/l)	260	-0.03 (-0.08, 0.01)	-0.04 (-0.09, 0.003) ^b
Log Cu (μg/l)			
Log Serum ferritin pre-pregnancy (µg/l)	258	-0.02 (-0.06, 0.01)	-0.03 (-0.06; 0.004) ^c
Log Serum ferritin at ANC2 (μg/l)	258	-0.03 (-0.07, 0.01)	-0.02 (-0.05; 0.02) ^c
Log Pb (µg/l)			
Log Serum ferritin pre-pregnancy (µg/l)	259	-0.04 (-0.10, 0.02)	-0.06 (-0.12, -0.006) ^d
Log Serum ferritin at ANC2 (μg/l)	260	-0.07 (-0.14, 0.001)	-0.11 (-0.17, -0.04) ^d
Log As (μg/l)			
Log Serum ferritin pre-pregnancy (µg/l)	201	-0.03 (-0.10, 0.04)	-0.02 (-0.09, 0.05) ^a
Log Serum ferritin at ANC2 (μg/l)	201	-0.07 (-0.15, 0.004)	-0.06 (-0.13, 0.02) ^a
Se (µg/l)			

Serum ferritin pre-pregnancy (µg/l)	201	0.16 (0.004, 0.31)	0.19 (0.043, 0.34) ^e
Serum ferritin at ANC2 (μg/l)	201	0.11 (0.010, 0.22)	0.11 (0.013, 0.22) ^e
Log Hg (μg/l)			
Log Serum ferritin pre-pregnancy (µg/l)	201	0.003 (0.000, 0.01)	-0.07 (-0.01, 0.16) ^f
Log Serum ferritin at ANC2 (μg/l)	201	0.002 (-0.000, 0.003)	0.03 (-0.06, 0.13) ^f

319 ^aAdjusted for G2 gestational age, gravidity, education, wealth and time from pre-pregnancy blood test to conception

320 ^bAdjusted for G2 gestational age, gravidity, education and wealth

321 cAdjusted for G2 gestational age, gravidity, education, wealth, time from pre-pregnancy blood test to conception, BMI and
 322 living lake-side

323 ^dAdjusted for G2 gestational age, gravidity, education, wealth and living lake-side

324 eAdjusted for G2 gestational age, gravidity, education, wealth and BMI

- 325 ^fAdjusted for G2 gestational age, gravidity, education, wealth, BMI and living lake-side
- 326

327 4. Discussion

328 These results show that serum ferritin concentrations prior to conception have an influence on blood

329 concentrations of iron, Mn, Cd, Pb and Se during pregnancy. Pre-pregnancy serum ferritin

- 330 concentrations were inversely associated with higher whole blood concentrations of Mn, Cd and Pb
- and positively associated with higher Se concentrations in the first trimester. With iron deficiency
- 332 pre-pregnancy strongly related to iron deficiency in the first trimester, similar associations are seen
- between ANC2 serum ferritin concentrations and Mn, Cd, Pb and Se, as would be expected.

334 4.1 Iron

The number of women found to be anemic prior to pregnancy was high (57.9%) with that due to iron

deficiency at 17%. Although iron deficiency is commonly assumed to cause half of all cases of

anemias (WHO, 2015), this figure is in line with a recent meta-analysis of national survey data which

found where anemia prevalence was greater than 40% in rural areas the percentage associated with

iron deficiency was 16% (Petry et al., 2016).

340 All women received iron supplements from their first ANC visit and overall serum ferritin

341 concentrations increased, and the numbers of anemic and/or iron deficient women decreased

342 between the pre-pregnancy and ANC2 visit blood tests implying that supplementation was effective.

343 It was not however possible to know from the data collected the compliance of individual women in

344 regularly taking iron supplements. Women were asked at each ANC visit if they had taken iron

345 supplements the previous day but no relationship was found between the number of positive

346 responses and iron status. Iron status prior to pregnancy had an important effect on iron status in

early pregnancy. The odds of being iron deficient at the ANC2 visit were 41.2 (95%Cl 6.2;275.0) times

- 348 more likely if iron deficient pre-pregnancy. The wide confidence interval of this result reflects the
- 349 small number of women in the sample who remained iron deficient at ANC2. For this reason, a
- 350 different cutoff was also tested based on an estimate of iron stores that are needed to be present at
- 351 the start of pregnancy to fully meet requirements and was consistent with the previous result with
- 352 the odds of a woman remaining without sufficient iron stores for pregnancy at ANC2 being 12.5
- 353 (95%Cl 5.11;30.6) times higher if low in iron stores prior to pregnancy. This finding suggests that for
- 354 women who are not classified as iron deficient but have lower iron stores, the iron supplementation
- that they receive is not sufficient to carry them through all the iron requirements of pregnancy.

356 The current recommendations for iron supplementation in pregnancy in Benin is daily oral iron and 357 folic acid supplementation usually provided through ANC visits and coverage is relatively high with 358 85% of women who had a live birth in the previous 5 years reporting having taken oral iron 359 supplement in the most recent national demographic and health survey (INSAE & ICF, 2019). A 360 comparison of rates of adherence to iron supplementation during pregnancy between 22 sub-Saharan found Benin had the 4th highest reported rate of these countries and varies as may be 361 expected with the number of antenatal visits, and levels of education and family wealth (Ba et al., 362 363 2019). It appears that iron supplementation is well accepted in Benin but may start later than ideal as 364 seen in some of the quite late first ANC visits in this cohort, despite regular follow-up. It is also 365 possible that the same socio-economic factors that contribute to lower adherence to iron 366 supplementation may also contribute to higher environmental exposures to metals.

367 4.2 Manganese, Cadmium and Lead

Blood Mn concentrations in the range 4–15 µg/L are considered normal (ATSDR, 2012b). Although 368 369 the majority of blood samples taken at ANC2 in this study fell within this range, 15% of women 370 recorded concentrations above 15 µg/L. As blood manganese concentrations are known to increase 371 during pregnancy and be significantly higher in cord than in maternal blood samples at delivery 372 (Gunier et al., 2014; Takser et al., 2004; Tholin et al., 1995; Yamamoto et al., 2019), it is likely that an 373 even greater percentage of women would have been exposed to higher concentrations of Mn by the 374 end of pregnancy than that measured at ANC2. Several studies link adverse effects on birth size 375 parameters with both low and high concentrations of Manganese in maternal and/or cord blood 376 (Guan et al., 2014; Yamamoto et al., 2019; Zota et al., 2009) and between higher concentrations of 377 manganese and impaired child neurodevelopment (Chung et al., 2015; Claus Henn et al., 2017; Lin et 378 al., 2013; Takser et al., 2003). Evidence from other studies also supports a close interaction between 379 manganese and iron, particularly in an iron deficient state, in terms of their transport and absorption 380 mechanisms into the body (Bjørklund et al., 2017; Margrete Meltzer et al., 2010). Blood Mn 381 concentrations have been found to be higher among pregnant women (Tholin et al., 1995). Low iron 382 status may also increase manganese concentration in certain brain areas as demonstrated in animal 383 studies (Erikson et al., 2004, 2005).

384 The results for Mn, Cd and Pb appear to be consistent with reported transport affinities of DMT1 whereby Mn and Cd are transported in preference to iron and Pb (Garrick et al., 2006). The slightly 385 386 lower increase in Pb concentrations compared to Mn and Cd for the same change in serum ferritin 387 concentrations for both pre-pregnancy and ANC2 results may indicate the competition between 388 these metals for absorption through this transport mechanism. A similar pattern was found in a Norwegian cohort monitored for metals mid-pregnancy where blood concentrations of Mn and Cd 389 390 were found to be significantly higher in women who were iron deficient, with a greater difference in 391 Mn concentrations compared to Cd concentrations. Pb concentrations in this cohort however were 392 not found to be significantly different between iron deficient and non-iron deficient women 393 (Caspersen et al., 2019).

394 4.3 Selenium

395 For optimal activity of Se dependent proteins, concentrations in whole blood would be expected to 396 be in the range 125-163 μ g/L (Muecke et al., 2018). This is similar to the concentrations measured in 397 the first trimester for women in this study where the median value was 149.1 and the inter-quartile range 132.1-167.3 µg/L. Although a significant positive association was found between serum ferritin 398 399 concentrations and Se concentrations, the effect was guite small. For example all other parameters 400 being equal, the difference between Se concentration if increasing from the minimum pre-pregnancy 401 serum ferritin concentration of 4.3 to the average of 42 µg/l would be an increase of 7 µg/l which is a 402 relatively small difference on the scale of the Se results measured. Similar results were found in a 403 mid-pregnancy cohort where a small but significant increase in blood Se concentrations was found in 404 non iron deficient women compared to iron deficient women (Caspersen et al., 2019). The adjusted R 405 squared values for the Se models in this study were the lowest of those that found a significant 406 association explaining only 7% to 8% of the variation in the dataset indicating that other important 407 factors are certainly influencing Se values. Although there is some evidence that low selenium 408 concentrations can be associated with anaemia and iron deficiency anaemia, increasing iron stores is 409 not thought to have a direct effect on selenium uptake (Gürgöze et al., 2004; Semba et al., 2009; 410 Viita et al., 1989; Yetgin et al., 1992).

411 4.4 Limitations and Strengths

412 There are several factors that add to measurement uncertainty in the dataset. The time between the 413 blood draw before pregnancy and pregnancy confirmation varied widely from just over 2 weeks to 414 almost 2 years and iron status may have changed over time. There were also differences in the 415 timing of ANC visits meaning that iron supplementation would have started at different points during 416 the pregnancy and that what was intended as the first trimester blood draw did not take place until 417 early in the second trimester for 16% of women in the sample and both of these could have affected 418 the concentrations of metals measured. In particular Mn has a relatively low half-life in blood of 419 around 30 days (Finley et al., 2003) so changes in iron status between the pre-pregnancy and ANC2 420 blood draw may not fully reflect Mn exposure over that time period. Nevertheless, there was a 421 strong correlation between serum ferritin concentrations at the pre-pregnancy and ANC2 blood 422 draws (r_s=0.65) which implies that the differing time periods did not change relationships too much. 423 If some women experienced longer periods of iron supplementation than others, this may have 424 resulted in some low pre-pregnancy serum ferritin concentrations not corresponding to higher 425 concentrations of Mn at ANC2. This would have reduced the strength of the associations found in the 426 regression analysis as well as reducing the value of the coefficients. Conversely, for the women who 427 were not tested for metals until the second trimester of pregnancy, metal concentrations may look 428 higher than they actually were for a normal serum ferritin concentration, given that the uptake of 429 some metals increase during pregnancy. This would have had the opposite effect on the regression 430 coefficients, potentially increasing above their true value while also reducing the strength of the 431 association.

The major strength of this study is that detailed information was collected prior to conception which
is not often available in comparable pregnancy cohorts. The number of metals assessed and the
ability to correct for inflammation in this context adds to understanding of environmental exposures

- in Benin for which there are few existing studies. It highlights that women with lower iron stores
- 436 prior to pregnancy, even if not classified as iron deficient, are at risk of absorbing higher amounts of
- 437 metals from their environments that are detrimental to the health of both mother and child. It is
- 438 suggested that testing the effects of iron and folic acid supplementation beginning prior to
- 439 conception on concentrations of Mn, Cd and Pb during pregnancy would be worthwhile to determine
- 440 if fetal exposures can be reduced without any other negative effects.

441 5. Conclusion

- 442 This study shows that serum ferritin concentrations are negatively associated with increased uptake
- of the toxic metals Mn, Cd and Pb in pregnancy. Importantly, lower serum ferritin concentrations
- 444 prior to conception were found to be associated with higher concentrations of Mn, Cd and Pb
- suggesting that increasing iron both before and during pregnancy may limit uptake of these metals.
- 446 These results argue in favour of testing the effects of iron supplementation prior to pregnancy to
- 447 better safeguard foetal development.

448 **6. References**

- 449 Accrombessi, M., Yovo, E., Cottrell, G., Agbota, G., Gartner, A., Martin-Prevel, Y., Fanou-Fogny, N.,
- 450 Djossinou, D., Zeitlin, J., Tuikue-Ndam, N., Bodeau-Livinec, F., Houzé, S., Jackson, N.,
- 451 Ayemonna, P., Massougbodji, A., Cot, M., Fievet, N., & Briand, V. (2018). Cohort profile :
- 452 Effect of malaria in early pregnancy on fetal growth in Benin (RECIPAL preconceptional
- 453 cohort). *BMJ Open*, 8(1), e019014. https://doi.org/10.1136/bmjopen-2017-019014
- Anderson, G. J., & Frazer, D. M. (2017). Current understanding of iron homeostasis. *The American*Journal of Clinical Nutrition, 106(suppl_6), 1559S-1566S.
- 456 https://doi.org/10.3945/ajcn.117.155804
- 457 Arnaud, J., & van Dael, P. (2018). Selenium Interactions with Other Trace Elements, with Nutrients
- 458 (and Drugs) in Humans. In B. Michalke (Éd.), *Selenium* (p. 413-447). Springer International
- 459 Publishing. https://doi.org/10.1007/978-3-319-95390-8_22
- 460 ATSDR. (2012a). *Toxicological Profile for Cadmium*. 487.
- 461 ATSDR. (2012b). Toxicological Profile for Manganese. U.S. Department of Health and Human Services,
 462 Public Health Service Agency for Toxic Substances and Disease Registry, 556.
- 463 ATSDR Toxicological Profile : Lead. (s. d.). Consulté 20 août 2020, à l'adresse

464 https://www.atsdr.cdc.gov/toxprofiles/tp.asp?id=96&tid=22

- 465 Ba, D. M., Ssentongo, P., Kjerulff, K. H., Na, M., Liu, G., Gao, X., & Du, P. (2019). Adherence to Iron
- 466 Supplementation in 22 Sub-Saharan African Countries and Associated Factors among
- 467 Pregnant Women : A Large Population-Based Study. *Current Developments in Nutrition*,
- 468 *3*(12). https://doi.org/10.1093/cdn/nzz120
- 469 Bellinger, D. C. (2005). Teratogen update : Lead and pregnancy. Birth Defects Research. Part A,
- 470 Clinical and Molecular Teratology, 73(6), 409-420. https://doi.org/10.1002/bdra.20127
- 471 Bjørklund, G., Aaseth, J., Skalny, A. V., Suliburska, J., Skalnaya, M. G., Nikonorov, A. A., & Tinkov, A. A.
- 472 (2017). Interactions of iron with manganese, zinc, chromium, and selenium as related to
- 473 prophylaxis and treatment of iron deficiency. Journal of Trace Elements in Medicine and

474	Biology: Organ of the Society for Minerals and Trace Elements (GMS), 41, 41-53.
475	https://doi.org/10.1016/j.jtemb.2017.02.005
476	Bothwell, T. H. (2000). Iron requirements in pregnancy and strategies to meet them. The American
477	Journal of Clinical Nutrition, 72(1), 257S-264S. https://doi.org/10.1093/ajcn/72.1.257S
478	Bressler. (s. d.). Divalent Metal Transporter 1 in Lead and Cadmium Transport—BRESSLER - 2004—
479	Annals of the New York Academy of Sciences—Wiley Online Library. Consulté 14 août 2019, à
480	l'adresse
481	https://nyaspubs.onlinelibrary.wiley.com/doi/abs/10.1196/annals.1306.011?sid=nlm%3Apu
482	bmed
483	Budtz-Jørgensen, E., Bellinger, D., Lanphear, B., Grandjean, P., & International Pooled Lead Study
484	Investigators. (2013). An international pooled analysis for obtaining a benchmark dose for
485	environmental lead exposure in children. Risk Analysis: An Official Publication of the Society
486	for Risk Analysis, 33(3), 450-461. https://doi.org/10.1111/j.1539-6924.2012.01882.x
487	Caspersen, I. H., Thomsen, C., Haug, L. S., Knutsen, H. K., Brantsæter, A. L., Papadopoulou, E., Erlund,
488	I., Lundh, T., Alexander, J., & Meltzer, H. M. (2019). Patterns and dietary determinants of
489	essential and toxic elements in blood measured in mid-pregnancy : The Norwegian
490	Environmental Biobank. Science of The Total Environment, 671, 299-308.
491	https://doi.org/10.1016/j.scitotenv.2019.03.291
492	CDC. (2010). Leadandpregnancy2010.pdf.
493	https://www.cdc.gov/nceh/lead/publications/leadandpregnancy2010.pdf
494	Chung, S. E., Cheong, HK., Ha, EH., Kim, BN., Ha, M., Kim, Y., Hong, YC., Park, H., & Oh, SY.
495	(2015). Maternal Blood Manganese and Early Neurodevelopment : The Mothers and
496	Children's Environmental Health (MOCEH) Study. Environmental Health Perspectives, 123(7),
497	717-722. https://doi.org/10.1289/ehp.1307865
498	Claus Henn, B., Bellinger, D. C., Hopkins, M. R., Coull, B. A., Ettinger, A. S., Jim, R., Hatley, E.,
499	Christiani, D. C., & Wright, R. O. (2017). Maternal and Cord Blood Manganese Concentrations

500	and Early Childhood Neurodevelopment among Residents near a Mining-Impacted Superfund
501	Site. Environmental Health Perspectives, 125(6). https://doi.org/10.1289/EHP925
502	Erikson, K. M., Syversen, T., Aschner, J. L., & Aschner, M. (2005). Interactions between excessive
503	manganese exposures and dietary iron-deficiency in neurodegeneration. Environmental
504	<i>Toxicology and Pharmacology, 19</i> (3), 415-421. https://doi.org/10.1016/j.etap.2004.12.053
505	Erikson, K. M., Syversen, T., Steinnes, E., & Aschner, M. (2004). Globus pallidus : A target brain region
506	for divalent metal accumulation associated with dietary iron deficiency. The Journal of
507	Nutritional Biochemistry, 15(6), 335-341. https://doi.org/10.1016/j.jnutbio.2003.12.006
508	Esteban-Vasallo María D., Aragonés Nuria, Pollan Marina, López-Abente Gonzalo, & Perez-Gomez
509	Beatriz. (2012). Mercury, Cadmium, and Lead Levels in Human Placenta : A Systematic
510	Review. Environmental Health Perspectives, 120(10), 1369-1377.
511	https://doi.org/10.1289/ehp.1204952
512	Finley, J. W., Penland, J. G., Pettit, R. E., & Davis, C. D. (2003). Dietary Manganese Intake and Type of
513	Lipid Do Not Affect Clinical or Neuropsychological Measures in Healthy Young Women. The
514	Journal of Nutrition, 133(9), 2849-2856. https://doi.org/10.1093/jn/133.9.2849
515	Garrick, M. D., Singleton, S. T., Vargas, F., Kuo, HC., Zhao, L., Knöpfel, M., Davidson, T., Costa, M.,
516	Paradkar, P., Roth, J. A., & Garrick, L. M. (2006). DMT1 : Which metals does it transport?
517	<i>Biological Research, 39</i> (1), 79-85. https://doi.org/10.4067/S0716-97602006000100009
518	Georgieff, M. K., Ramel, S. E., & Cusick, S. E. (2018). Nutritional influences on brain development.
519	Acta Paediatrica, 107(8), 1310-1321. https://doi.org/10.1111/apa.14287
520	Guan, H., Wang, M., Li, X., Piao, F., Li, Q., Xu, L., Kitamura, F., & Yokoyama, K. (2014). Manganese
521	concentrations in maternal and umbilical cord blood : Related to birth size and
522	environmental factors. European Journal of Public Health, 24(1), 150-157.
523	https://doi.org/10.1093/eurpub/ckt033

- 524 Gundacker, C., & Hengstschläger, M. (2012). The role of the placenta in fetal exposure to heavy
- 525 metals. *Wiener Medizinische Wochenschrift*, *162*(9), 201-206.

526 https://doi.org/10.1007/s10354-012-0074-3

- 527 Gunier, R. B., Mora, A. M., Smith, D., Arora, M., Austin, C., Eskenazi, B., & Bradman, A. (2014).
- 528 Biomarkers of Manganese Exposure in Pregnant Women and Children Living in an
- 529 Agricultural Community in California. *Environmental Science & Technology*, 48(24),

530 14695-14702. https://doi.org/10.1021/es503866a

- 531 Gürgöze, M. K., Denizmen Aygün, A., Ölçücü, A., Doğan, Y., & Yılmaz, E. (2004). Plasma selenium
- 532 status in children with iron deficiency anemia. *Journal of Trace Elements in Medicine and*

533 *Biology*, *18*(2), 193-196. https://doi.org/10.1016/j.jtemb.2004.07.004

- 534 Guy, M., Accrombessi, M., Fievet, N., Yovo, E., Massougbodji, A., Le Bot, B., Glorennec, P., Bodeau-
- 535 Livinec, F., & Briand, V. (2018). Toxics (Pb, Cd) and trace elements (Zn, Cu, Mn) in women
- 536 during pregnancy and at delivery, South Benin, 2014–2015. Environmental Research, 167,
- 537 198-206. https://doi.org/10.1016/j.envres.2018.06.054
- 538 INSAE, & ICF. (2019). Enquête Démographique et de Santé au Bénin, 2017-2018. Institut National de
- 539 la Statistique et de l'Analyse Économique (INSAE) et ICF.
- 540 https://dhsprogram.com/pubs/pdf/FR350/FR350.pdf
- Kim, Y. (2018). Effect of Iron Deficiency on the Increased Blood Divalent Metal Concentrations. *Iron Deficiency Anemia*. https://doi.org/10.5772/intechopen.78958
- 543 Kim, Y., Kim, B.-N., Hong, Y.-C., Shin, M.-S., Yoo, H.-J., Kim, J.-W., Bhang, S.-Y., & Cho, S.-C. (2009). Co-
- 544 exposure to environmental lead and manganese affects the intelligence of school-aged
- 545 children. *Neurotoxicology*, *30*(4), 564-571. https://doi.org/10.1016/j.neuro.2009.03.012
- 546 Kippler, M., Tofail, F., Hamadani, J. D., Gardner, R. M., Grantham-McGregor, S. M., Bottai, M., &
- 547 Vahter, M. (2012). Early-Life Cadmium Exposure and Child Development in 5-Year-Old Girls
- 548 and Boys : A Cohort Study in Rural Bangladesh. *Environmental Health Perspectives*, 120(10),
- 549 1462-1468. https://doi.org/10.1289/ehp.1104431

- 550 Lin, C.-C., Chen, Y.-C., Su, F.-C., Lin, C.-M., Liao, H.-F., Hwang, Y.-H., Hsieh, W.-S., Jeng, S.-F., Su, Y.-N.,
- 551 & Chen, P.-C. (2013). In utero exposure to environmental lead and manganese and
- neurodevelopment at 2 years of age. *Environmental Research*, *123*, 52-57.
- 553 https://doi.org/10.1016/j.envres.2013.03.003
- Loréal, O., Cavey, T., Bardou-Jacquet, E., Guggenbuhl, P., Ropert, M., & Brissot, P. (2014). Iron,
- hepcidin, and the metal connection. *Frontiers in Pharmacology*, 5.
- 556 https://doi.org/10.3389/fphar.2014.00128
- 557 Madejczyk, M. S., & Ballatori, N. (2012). The iron transporter ferroportin can also function as a
- 558 manganese exporter. *Biochimica Et Biophysica Acta*, *1818*(3), 651-657.
- 559 https://doi.org/10.1016/j.bbamem.2011.12.002
- 560 Mandy, M., & Nyirenda, M. (2018). Developmental Origins of Health and Disease : The relevance to
- 561 developing nations. *International Health*, *10*(2), 66-70.
- 562 https://doi.org/10.1093/inthealth/ihy006
- 563 Margrete Meltzer, H., Lise Brantsæter, A., Borch-Johnsen, B., Ellingsen, D. G., Alexander, J.,
- 564 Thomassen, Y., Stigum, H., & Ydersbond, T. A. (2010). Low iron stores are related to higher
- 565 blood concentrations of manganese, cobalt and cadmium in non-smoking, Norwegian
- 566 women in the HUNT 2 study. *Environmental Research*, *110*(5), 497-504.
- 567 https://doi.org/10.1016/j.envres.2010.03.006
- 568 Menezes-Filho, J. A., Carvalho, C. F., Rodrigues, J. L. G., Araújo, C. F. S., dos Santos, N. R., Lima, C. S.,
- 569 Bandeira, M. J., Marques, B. L. de S., Anjos, A. L. S., Bah, H. A. F., Abreu, N., Philibert, A., &
- 570 Mergler, D. (2018). Environmental Co-Exposure to Lead and Manganese and Intellectual
- 571 Deficit in School-Aged Children. International Journal of Environmental Research and Public
- 572 *Health*, *15*(11). https://doi.org/10.3390/ijerph15112418
- 573 Moya-Alvarez, V., Cottrell, G., Ouédraogo, S., Accrombessi, M., Massougbodgi, A., & Cot, M. (2017).
- 574 High Iron Levels Are Associated with Increased Malaria Risk in Infants during the First Year of

- 575 Life in Benin. *The American Journal of Tropical Medicine and Hygiene*, *97*(2), 497-503.
- 576 https://doi.org/10.4269/ajtmh.16-0001
- 577 Muecke, R., Waldschock, K., Schomburg, L., Micke, O., Buentzel, J., Kisters, K., Adamietz, I. A., &
- 578 Huebner, J. (2018). Whole Blood Selenium Levels and Selenium Supplementation in Patients
- 579 Treated in a Family Doctor Practice in Golßen (State of Brandenburg, Germany) : A
- 580 Laboratory Study. *Integrative Cancer Therapies*, *17*(4), 1132-1136.
- 581 https://doi.org/10.1177/1534735418807971
- 582 Muriuki, J. M., Mentzer, A. J., Kimita, W., Ndungu, F. M., Macharia, A. W., Webb, E. L., Lule, S. A.,
- 583 Morovat, A., Hill, A. V. S., Bejon, P., Elliott, A. M., Williams, T. N., & Atkinson, S. H. (2019).
- 584 Iron Status and Associated Malaria Risk Among African Children. *Clinical Infectious Diseases:*
- 585 An Official Publication of the Infectious Diseases Society of America, 68(11), 1807-1814.
- 586 https://doi.org/10.1093/cid/ciy791
- 587 Mwangi, M. N., Echoka, E., Knijff, M., Kaduka, L., Werema, B. G., Kinya, F. M., Mutisya, R., Muniu, E.
- 588 M., Demir, A. Y., Verhoef, H., & Bourdet-Sicard, R. (2019). Iron Status of Kenyan Pregnant
- 589 Women after Adjusting for Inflammation Using BRINDA Regression Analysis and Other
- 590 Correction Methods. *Nutrients*, *11*(2). https://doi.org/10.3390/nu11020420
- 591 Namaste, S. M., Rohner, F., Huang, J., Bhushan, N. L., Flores-Ayala, R., Kupka, R., Mei, Z., Rawat, R.,
- 592 Williams, A. M., Raiten, D. J., Northrop-Clewes, C. A., & Suchdev, P. S. (2017). Adjusting
- 593 ferritin concentrations for inflammation : Biomarkers Reflecting Inflammation and Nutritional
- 594 Determinants of Anemia (BRINDA) project. The American Journal of Clinical Nutrition,
- 595 *106*(Suppl 1), 359S-371S. https://doi.org/10.3945/ajcn.116.141762
- 596 Petry, N., Olofin, I., Hurrell, R. F., Boy, E., Wirth, J. P., Moursi, M., Donahue Angel, M., & Rohner, F.
- 597 (2016). The Proportion of Anemia Associated with Iron Deficiency in Low, Medium, and High
- 598 Human Development Index Countries : A Systematic Analysis of National Surveys. *Nutrients*,
- 599 8(11). https://doi.org/10.3390/nu8110693

- 600 Radlowski, E. C., & Johnson, R. W. (2013). Perinatal iron deficiency and neurocognitive development.
- 601 Frontiers in Human Neuroscience, 7. https://doi.org/10.3389/fnhum.2013.00585
- Rahman, A., Kumarathasan, P., & Gomes, J. (2016). Infant and mother related outcomes from
- 603 exposure to metals with endocrine disrupting properties during pregnancy. *The Science of*
- 604 the Total Environment, 569-570, 1022-1031. https://doi.org/10.1016/j.scitotenv.2016.06.134
- Rossipal, E., Krachler, M., Li, F., & Micetic-Turk, D. (2000). Investigation of the transport of trace
- elements across barriers in humans : Studies of placental and mammary transfer. *Acta Paediatrica (Oslo, Norway: 1992), 89*(10), 1190-1195.
- 608 Sangaré, L., van Eijk, A. M., ter Kuile, F. O., Walson, J., & Stergachis, A. (2014). The Association
- 609 between Malaria and Iron Status or Supplementation in Pregnancy : A Systematic Review and
- 610 Meta-Analysis. *PLoS ONE*, *9*(2). https://doi.org/10.1371/journal.pone.0087743
- 611 Semba, R., Ricks, M., Ferrucci, L., Xue, Q.-L., Guralnik, J., & Fried, L. (2009). Low serum selenium is
- associated with anemia among older adults in the United States. *European journal of clinical nutrition*, 63(1), 93-99. https://doi.org/10.1038/sj.ejcn.1602889
- 614 Stephenson, J., Heslehurst, N., Hall, J., Schoenaker, D. A. J. M., Hutchinson, J., Cade, J., Poston, L.,
- 615 Barrett, G., Crozier, S., Kumaran, K., Yanjik, C., Barker, M., Baird, J., & Mishra, G. (2018).
- 616 Before the beginning : Nutrition and lifestyle in the preconception period and its importance
- 617 for future health. *Lancet (London, England), 391*(10132), 1830-1841.
- 618 https://doi.org/10.1016/S0140-6736(18)30311-8
- Sun, H.-J., Rathinasabapathi, B., Wu, B., Luo, J., Pu, L.-P., & Ma, L. Q. (2014). Arsenic and selenium
- 620 toxicity and their interactive effects in humans. *Environment International, 69*, 148-158.
- 621 https://doi.org/10.1016/j.envint.2014.04.019
- Takser, L., Lafond, J., Bouchard, M., St-Amour, G., & Mergler, D. (2004). Manganese levels during
- 623 pregnancy and at birth : Relation to environmental factors and smoking in a Southwest
- 624 Quebec population. *Environmental Research*, *95*(2), 119-125.
- 625 https://doi.org/10.1016/j.envres.2003.11.002

Metabolite Levels at Birth, and Child Psychomotor Development. NeuroToxicology, 24(4),
667-674. https://doi.org/10.1016/S0161-813X(03)00058-5
Tholin, K., Sandström, B., Palm, R., & Hallmans, G. (1995). Changes in Blood Manganese Levels During
Pregnancy in Iron Supplemented and non Supplemented Women. Journal of Trace Elements
in Medicine and Biology, 9(1), 13-17. https://doi.org/10.1016/S0946-672X(11)80003-9
Truska, P., Rosival, L., Balázová, G., Hinst, J., Rippel, A., Palusová, O., & Grunt, J. (1989). Blood and
placental concentrations of cadmium, lead, and mercury in mothers and their newborns.
Journal of Hygiene, Epidemiology, Microbiology, and Immunology, 33(2), 141-147.
Viita, L. M., Mutanen, M. L., & Mykkänen, H. M. (1989). Selenium-iron interaction in young women
with low selenium status. Journal of Human Nutrition and Dietetics, 2(1), 39-42.
https://doi.org/10.1111/j.1365-277X.1989.tb00006.x
WHO. (2015). The Global Prevalence of Anemia in 2011.
https://apps.who.int/iris/bitstream/handle/10665/177094/9789241564960_eng.pdf?sequen
ce=1
WHO. (2016). Prevalence of anaemia in women of reproductive age—Estimates by WHO region.
WHO; World Health Organization.
https://apps.who.int/gho/data/view.main.ANAEMIAWOMENPREVANEMIAREG?lang=en
WHO. (2020). WHO Guideline on use of ferritin concentrations to assess iron status in individuals and
populations. World Health Organization, Geneva.
https://www.who.int/publications/i/item/9789240000124
WHO recommendations on antenatal care for a positive pregnancy experience. (2016). World Health
Organization.
Organization.

Takser, L., Mergler, D., Hellier, G., Sahuquillo, J., & Huel, G. (2003). Manganese, Monoamine

626

- 649 Yamamoto, M., Sakurai, K., Eguchi, A., Yamazaki, S., Nakayama, S. F., Isobe, T., Takeuchi, A., Sato, T.,
- Hata, A., Mori, C., Nitta, H., Ohya, Y., Kishi, R., Yaegashi, N., Hashimoto, K., Mori, C., Ito, S.,
- 651 Yamagata, Z., Inadera, H., ... Katoh, T. (2019). Association between blood manganese level

652	during pregnancy and birth size : The Japan environment and children's study (JECS).
653	Environmental Research, 172, 117-126. https://doi.org/10.1016/j.envres.2019.02.007
654	Yetgin, S., Hincal, F., Baçaran, N., & Ciliv, G. (1992). Serum Selenium Status in Children with Iron
655	Deficiency Anemia. Acta Haematologica, 88(4), 185-188. https://doi.org/10.1159/000204683
656	Zota, A. R., Ettinger, A. S., Bouchard, M., Amarasiriwardena, C. J., Schwartz, J., Hu, H., & Wright, R. O.
657	(2009). Maternal Blood Manganese Levels and Infant Birth Weight. Epidemiology
658	(Cambridge, Mass.), 20(3), 367-373. https://doi.org/10.1097/EDE.0b013e31819b93c0
659	