



**HAL**  
open science

## Calculation of Mercury's Effects on Neurodevelopment

Philippe Grandjean, Celine Pichery, Martine Bellanger, Esben  
Budtz-Jørgensen

► **To cite this version:**

Philippe Grandjean, Celine Pichery, Martine Bellanger, Esben Budtz-Jørgensen. Calculation of Mercury's Effects on Neurodevelopment. *Environmental Health Perspectives*, 2012, 120 (12), pp.A 452. 10.1289/ehp.1206033 . hal-03708463

**HAL Id: hal-03708463**

**<https://ehesp.hal.science/hal-03708463v1>**

Submitted on 29 Jun 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

The correspondence section is a public forum and, as such, is not peer-reviewed. EHP is not responsible for the accuracy, currency, or reliability of personal opinion expressed herein; it is the sole responsibility of the authors. EHP neither endorses nor disputes their published commentary.

## Calculation of Mercury's Effects on Neurodevelopment

<http://dx.doi.org/10.1289/ehp.1206033>

Bellinger (2012) recently estimated the loss of cognitive function in terms of Full-Scale intelligence quotient (IQ) in children exposed to certain environmental chemicals. To ascertain prenatal exposures of methylmercury (MeHg) in children, he used exposure data on mercury (Hg) concentrations in hair of U.S. women of childbearing age (16–49 years) from NHANES (National Health and Nutrition Examination Survey) 1999–2000 (McDowell et al. 2004). Bellinger applied a regression coefficient of  $-0.18$  IQ points per microgram per gram increase in maternal hair as calculated by Axelrad et al. (2007). However, the results of Axelrad et al. (2007) relied on incomplete data from a prospective study in the Faroe Islands and on non-adjusted results from the Seychelles study, later found to be confounded by nutrients from seafood (Strain et al. 2008). Bellinger (2012) then applied the regression coefficient to hair Hg levels  $> 1.11$   $\mu\text{g/g}$  (90th percentile), because this level corresponds to the reference dose of MeHg established many years ago. Assuming a concentration of  $1.73$   $\mu\text{g/g}$  (95th percentile) as the midpoint (rather than the average, which is higher) for the hair Hg levels of the 10% of U.S. women with a level  $> 1.11$   $\mu\text{g/g}$ , he estimated a total IQ loss of 284,580 points. We believe that Bellinger's general approach is sound but that the dose–response information is outdated, a caveat that Bellinger noted, although it was not reflected in the summary table. We therefore wish to complement these calculations using updated dose–response data.

Prospective data justify a lower threshold Hg level of  $0.58$   $\mu\text{g/g}$  hair corresponding to 50% of the reference dose (Grandjean and Budtz-Jørgensen 2007). In addition, a  $1$ - $\mu\text{g/g}$  increase in hair Hg concentration is more likely associated with an average adverse impact of  $0.465$  IQ points, as discussed by Pichery et al. (2012). Assuming a log-normal exposure distribution, a 75th percentile hair Hg concentration of  $0.42$   $\mu\text{g/g}$ , and a 90th percentile of  $1.11$   $\mu\text{g/g}$  as reported by McDowell et al. (2004), we estimate that 18.5% of women exceed a threshold of  $0.58$   $\mu\text{g/g}$  hair Hg and that the average concentration for  $0.58$ – $1.11$   $\mu\text{g/g}$  is approximately  $0.8$   $\mu\text{g/g}$ . For the sake of comparing these values with Bellinger's calculations (Bellinger 2012), we used a median concentration of  $1.73$   $\mu\text{g/g}$  as the average hair Hg

level of the 10% of U.S. women with a level  $> 1.11$   $\mu\text{g/g}$ . On the basis of these assumptions, we calculated a total IQ loss for the U.S. population of children 0–5 years of age ( $n = 25.5$  million) to be 1,590,000 IQ points, or 264,000 IQ points per year.

We recently used similar calculations to estimate the annual costs of Hg pollution in France (Pichery et al. 2012), a country one-fifth the size of the United States. At slightly higher exposure levels, the annual loss in IQ points was estimated to be 157,000. Greater losses were obtained using a log-scale effect (Pichery et al. 2012). With an estimated value of each IQ point of \$18,000 in terms of lifetime earnings, the current loss of IQ points associated with MeHg exposure represents a very substantial value to society.

*The authors declare they have no actual or potential competing financial interests.*

**Philippe Grandjean**

Harvard School of Public Health  
Boston, Massachusetts  
E-mail: pgrand@hsph.harvard.edu

**Celine Pichery**

**Martine Bellanger**  
School for Advanced Studies in  
Public Health (EHESP)  
Paris, France

**Esben Budtz-Jørgensen**

Department of Biostatistics  
University of Copenhagen  
Copenhagen, Denmark

### REFERENCES

- Axelrad DA, Bellinger DC, Ryan LM, Woodruff TJ. 2007. Dose–response relationship of prenatal mercury exposure and IQ: an integrative analysis of epidemiologic data. *Environ Health Perspect* 115:609–615.
- Bellinger DC. 2012. A strategy for comparing the contributions of environmental chemicals and other risk factors to neurodevelopment of children. *Environ Health Perspect* 120:501–507.
- Grandjean P, Budtz-Jørgensen E. 2007. Total imprecision of exposure biomarkers: implications for calculating exposure limits. *Am J Industr Med* 50:712–719.
- McDowell MA, Dillon CF, Osterloh J, Bolger PM, Pellizzari E, Fernando R, et al. 2004. Hair mercury levels in U.S. children and women of childbearing age: reference range data from NHANES 1999–2000. *Environ Health Perspect* 112:1165–1171.
- Pichery C, Bellanger M, Zmirou-Navier D, Fréry N, Cordier S, Roue-LeGall A, et al. 2012. Economic evaluation of health consequences of prenatal methylmercury exposure in France. *Environ Health* 11:53; doi:10.1186/1476-069X-11-53 [Online 10 August 2012].
- Strain JJ, Davidson PW, Bonham MP, Duffy EM, Stokes-Riner A, Thurston SW, et al. 2008. Associations of maternal long-chain polyunsaturated fatty acids, methyl mercury, and infant development in the Seychelles Child Development Nutrition Study. *Neurotoxicology* 29:776–782.

## Calculation of Mercury's Effects on Neurodevelopment: Bellinger Responds

<http://dx.doi.org/10.1289/ehp.1206033R>

In my paper (Bellinger 2012), I noted among the limitations that the calculations are only as valid as the data on which they are based. My hope was that those with a special interest in a particular risk factor would be stimulated to provide stronger data on either the exposure distribution or the dose–response relationship so that the calculations could be refined. I am therefore grateful to Grandjean et al. for providing an updated estimate of the dose–response relationship for prenatal methylmercury, the use of which suggests that the total Full-Scale IQ loss among U.S. children is considerably larger than my initial estimate. All of the estimates listed in Table 2 of my paper (Bellinger 2012) should be considered provisional and should be updated when more precise data become available.

*The author has served as an expert witness in civil litigation involving exposures of children to lead and metallic mercury and has received travel funding and honoraria to present lectures on environmental health of children.*

**David C. Bellinger**

Harvard Medical School  
Harvard School of Public Health  
Boston Children's Hospital  
E-mail: david.bellinger@childrens.harvard.edu

### REFERENCE

- Bellinger DC. 2012. A strategy for comparing the contributions of environmental chemicals and other risk factors to neurodevelopment of children. *Environ Health Perspect* 120:501–507.

## Estimating Risk of Circulatory Disease from Exposure to Low-Level Ionizing Radiation

<http://dx.doi.org/10.1289/ehp.1206046>

The comprehensive meta-analysis of Little et al. (2012) summarized possible circulatory disease risks related to medium and low doses of whole-body radiation exposure in humans. The authors looked at excess relative risk (ERR) estimates from 10 different epidemiological studies. Using two statistical measures to calculate pooled ERR, they determined aggregate measures of ERR for four detrimental health outcomes and they reported mostly significant values for the ERR per unit dose in their Table 2.

Nine of the 10 studies Little et al. (2012) considered included moderate cumulative doses  $> 0.4$  Sv (see their Table 1), and they observed that risk trends in most cohorts were driven by a relatively small number of