

Work environment differences between outpatient and inpatient surgery: a pilot study on the vascular surgeons' perceptions

Laurence Destrieux, Youssra Yemmas, Sydney Williams, Nolwenn Le Meur

▶ To cite this version:

Laurence Destrieux, Youssra Yemmas, Sydney Williams, Nolwenn Le Meur. Work environment differences between outpatient and inpatient surgery: a pilot study on the vascular surgeons' perceptions. Annals of Vascular Surgery, 2024, 104, pp.156-165. 10.1016/j.avsg.2023.12.080. hal-04517733

HAL Id: hal-04517733 https://ehesp.hal.science/hal-04517733

Submitted on 3 Jun 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



1 2	Work environment differences between outpatient and inpatient surgery: a pilot study on the vascular surgeons' perceptions
3	short title: Vascular surgeons' view of their work environment
4	Laurence Destrieux, MD¹, Youssra Yemmas², Sydney Williams³, Nolwenn Le Meur, PhD⁴,5
5	
6	Affiliations:
7 8	¹ Clinique Générale, Groupe Vivalto, Département de chirurgie vasculaire, 4 Chemin Tour La Reine, 74000 Annecy, France
9	² Department of Biostatistics, Columbia University, 116 th and Broadway, New York, NY 10027, USA
LO	³ Columbia University Vagelos College of Physicians and Surgeons, 630 W 168th St, New York, NY
l1	10032, USA
12	⁴ Univ Rennes, EHESP, CNRS, Inserm, Arènes-UMR 6051, RSMS-U 1309, F-35000 Rennes, France
13 14	⁵ Ecoles Des Hautes Etudes en Santé Publique, Département METIS, 15 Avenue du Professeur Léon Bernard, CS 74312, 35043 Rennes Cedex, France
15	
16	Corresponding author:
L7	Nolwenn Le Meur
18	Nolwenn.lemeur@ehesp.fr
19	Permanent address:
20	Ecoles Des Hautes Etudes en Santé Publique, Département METIS, 15 Avenue du Professeur Léon Bernard, CS
21	74312, 35043 Rennes Cedex, France
22	
23	
24	
25	
26	
27	
28	
29	
30	

Abstract

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

Background: In France, outpatient endovascular surgical procedures have been slowly implemented in hospitals since 2015. Their development has been heterogeneous across France and is not yet the standard of care despite their benefits concerning patients' outcomes and healthcare professionals' satisfaction. However, since the COVID-19 pandemic, the demand for patient beds has been increasing, while human resources have been decreasing. This encouraged the surgery service reorganization and accelerated the shift from inpatient to outpatient surgery. Consequently, services had to adapt rapidly and this may have caused some strain on the hospital medical workforce. The objectives of this pilot study were to document and analyze the nursing staff (nurses and certified assistant nurses) well-being and workload as perceived by the vascular surgeons working with them. It also wanted to assess the link between the nursing staff's psychosocial stress and the surgeons' concentration and serenity in their workplace, based on the assumption that the well-being of both parties is inextricably linked. Methods: An observational study was conducted using an online questionnaire distributed to the senior members of the French society of vascular and endovascular surgery (n=490) between October 10 and October 18, 2022. Results: In total, 125 surgeons completed the questionnaire (25% response rate). The main finding was that according to 68% of vascular surgeons, the nursing staff's psychosocial stress significantly impaired their serenity and concentration at work and this frequently affected the surgical procedure safety. The main sources of psychosocial stress were the high work pace (64%), demand for flexibility (56%), lack of anticipation of schedule changes (82%), and difficulties encountered in relaying these concerns to hospital managers (44.6%). Conclusions: This study demonstrated that concomitantly with the forced acceleration of outpatient activity implementation, the vascular surgeons' perceptions of their working environment is deteriorating, especially

in conventional (inpatient) surgery wards where the workload is increasing and patients have more

comorbidities. The worsened psychosocial stress of surgeons and staff affects the care provided.

Introduction

56

57

58

59

60

61

62

63

64

65

66

67

68

69

70

71

72

73

74

75

76

77

78

79

80

81

In France, a country with an aging population, high tobacco consumption and modern lifestyle, the increasing needs in vascular surgery specialists was foreseen by the French College of Vascular Surgery and healthcare services should have been progressively adapted (1,2). In addition, recent technical progress in minimally invasive surgery for the treatement of peripheral aterial diseases should have contributed to the developmenent of same-day discharge endovascular revascularization and help with the organization of vascular care. However, in France, endovascular surgery procedures for peripheral artery diseases are mostly performed in inpatient settings (i.e. conventional wards) (3). Inpatient surgery includes at least an overnight stay. Conversely, in outpatient surgery, patients are hospitalized less than 12 hours, without overnight stay (same-day discharge). Several studies demonstrated that outpatient surgery presents benefits in patient's outcomes and work organization (4–8). However, outpatient endovascular surgery procedures have been slowly implemented in French hospitals since 2015 (3), their development has been heterogeneous, and they are not yet the standard of care despite their documented benefits (4). Concurrently, the COVID-19 crisis overstretched healthcare services and imposed substantial adjustments in healthcare services (9,10). Notably in vascular services, it encouraged the reorganization of surgery wards and the move from inpatient to outpatient surgery (the so-called "ambulatory shift") (11,12). Since the COVID-19 pandemic, the ambulatory shift seems to have accelerated in France. However the COVID-19 crisis also accelerated the defection of healthcare professionals, thus increasing the burden of care, particularly in conventional vascular surgery wards where the demand for patient beds has been increasing. Therefore, the combine impact of the ambulatory shift and post-COVID-19 health care professional ressources on vascular surgery wards needs to be documented. A major effect concerns the human resource redistribution between conventional and day surgery wards that threatens healthcare planning (13), especially in a context of healthcare workforce shortage (14). In addition, several vascular surgeons raised concerns about the degradation of life at work and more globally on the psychological stress of healthcare workforces, particularly in conventional wards. Many studies demonstrated that the degradation of work life is associated with the degradation of the care quality and safety (15–18). However, only few reports focused on the healthcare workforces' psychological

stress and its impact on care quality and safety in the context of the shift from inpatient to outpatient care (19,20). Consequently, it becomes critical to better assess the quality of life at work in surgery wards in the context of the workforce shortage and the increase of outpatient surgery procedures because it may affect healthcare service organization and patient care.

The objective of this pilot study was to evaluate the impact of the ambulatory shift due to the COVID-19 pandemic on the vascular surgeons' quality of work, with a focus on conventional surgery wards, and their perception of the healthcare staff's psychological stress. Specifically, we addressed the following questions:

(1) How do vascular surgeons perceive the healthcare staff's quality of work life? (2) How does the healthcare staff's quality of work life affect the vascular surgeons' serenity and concentration? (3) How does the healthcare staff's quality of work life affect the safety of the vascular surgeons' procedures?

Methods

Study design and participants

An observational study was conducted from the 10 to the 18 October 2022. It consisted of a voluntary online survey sent to senior vascular surgeons, members of the French society of vascular and endovascular surgery.

Questions and variables

The questions focused on the working environment with an emphasis on the nursing staff (i.e. nurses and certified assistant nurses) working in vascular surgery wards. The survey consisted of 28 questions divided in six main topics: characteristics of the hospital or the clinic, impact of the COVID-19 pandemic on surgical activity, discrepancy between surgical activity and nursing resources, nursing staff's work conditions, impact of the nursing staff's work conditions on the vascular surgeons' work environment, and interpersonal relationships and communication between vascular surgeons and the human resources department. The

work environmental factors were described and analyzed in order to understand the vascular surgeons' answer to the following questions: (1) "In your opinion, are nursing staff unhappy with their working conditions (high levels of occupational stress, anxiety, unsatisfaction)?" [do not know/no, rarely/yes, most of the time] (2) "Are your serenity and professional concentration affected by the nursing staff's quality of work life (organization, skills, experiences)?" (3) "Is the safety of your surgical interventions affected by the nursing staff's quality of work life?" [exceptionally/occasionally/frequently/difficult to assess]. The complete questionnaire is available in Supplementary material.

Statistical analysis

Univariate statistical analyses were performed using Fisher's exact tests or Pearson's χ^2 tests. Variables with a p-value <0.25 were included in the multivariate models. Multivariate logistic regression models were aplied using stepwise approaches to analyze the effect of each variable on the questions of interest, all other things being equal. P-values < 0.05 were considered statistically significant. Data management and statistical analyses were performed using the R statistical software (version 4.1.2)

Results

Characteristics of the respondents

In total, 125 vascular surgeons completed the survey (25.5% of the 490 senior members of the French society of vascular and endovascular surgery). The sample was representative of distribution of the vascular surgeons' in the different hospital sectors: 53% in private-sector hospitals and 41% in public-sector hospitals. Most surgeons (86%) performed both arterial and venous surgical procedures. Although 40% of respondents said that the COVID-19 health crisis led to an increase in their outpatient activities, the proportion of outpatient arterial surgery procedures remained lower than 10% for 48% of respondents.

Nursing staff resources

Most surgical units operated with a 8-13 patients per nurse ratio, irrespective of the setting. Only 13% of respondents who worked in day surgery wards and 18% of those who worked in conventional surgery wards reported a patient-to-nurse ratio >13. It should be noted that 27% and 4% of respondents could not assess this ratio for their day surgery ward and conventional surgery ward, respectively. Overall, nursing staff in day surgery wards seemed to cover more medical and surgical specialties than in conventional wards (49% vs. 9.7%).

Factors associated with the nursing staff's quality of work life

129

130

131

132

133

134

135

136

137

138

139

140

141

142

143

144

145

146

147

148

149

150

151

152

According to 52% and 37% of respondents, overall, the nursing staff's workload has been increasing in conventional and day surgery wards, respectively. Whatever the ward, nursing staff needed to be flexible; however, 50% of responders highlighted a lack of information to anticipate changes. Most respondents (65%) underlined that in conventional wards, the work pace was very sustained and 50% of respondents indicated that from 15% to >30% of errors in nursing care were due to time constraints (23% in day surgery wards). Conversely, 42% of respondents considered the work pace to be adapted or even low in day surgery wards. However, it should be noted that 21% of respondents were unable to assess this point. Moreover, 57% of respondents said that the supervision of nursing staff was deficient and 50% found that discussion with hospital managers was difficult when psychological stress was observed in nursing staff members. Overall, 68% of respondents (n=85) reported frequent psychological stress in their nursing staff. Only two respondents said that nursing staff did not show any stress and five could not assess it. These response modalities were grouped with the "Sometimes" versus "Often" to maintain the statistical power in the multivariate analyses. When analyzed independently, changes in the conventional surgery activity linked to the COVID-19 pandemic, work pace, supervision, flexibility, anticipation of schedule changes, and interactions with hospital managers were significantly associated with frequent work-related psychosocial stress in nursing staff (Table I and II). In the multivariate analysis with all variables adjusted for each other, demand for flexibility, sustained work pace in conventional surgery wards, and late notice of planning

changes contributed significantly to work-related psychosocial stress in the nursing staff (Table III), but not perceived lack of supervision and patient-to-nurse ratio.

Factors associated with the vascular surgeons' quality of work life

Concerning the surgery procedure safety in conventional surgery wards, the vascular surgeons' perception of the impact of the nursing staff's quality of work life was not clear-cut: 53% and 28% mentioned occasional and negative effects, respectively (Table IV). The significant and independently contributing factors were the frequent psychosocial stress of nursing staff, increased complexity of patients, and sustained work pace.

Compared with surgeons who reported that nursing staff's issues only rarely affected the safety of their surgical practice, surgeons who said that such issues frequently perturbed their work safety also reported significantly more nursing staff's psychosocial stress and more patients with multiple chronic conditions (Table V). The odds ratio for the work pace suggested the possible negative effect of sustained pace of work on patient safety, but overall, this was not significant.

The factors significantly and independently associated with the vascular surgeons' reduced serenity and concentration were the patients' increased level of comorbidities, increased lack of nursing care due to time constraints, frequent psychosocial stress of nursing staff, and difficulties in discussing the nursing staff's working conditions with hospital managers (Table VI). Compared with responders who reported rare perturbations, vascular surgeons who reported frequent impairment of their serenity and concentration also reported 4 times more nursing staff's psychosocial stress and higher number of comorbidities in patients (Table VII). The absence of mediation with hospital managers also was significantly correlated with occasional and frequent perturbations compared with rare. A surgeon stated that "our serenity is no longer affected, but we lose a lot of energy in arguing to maintain sufficient workforce levels" and another added that "... we see more and more staff turnover, with loss of specific knowledge on vascular patients' management". These qualitative statements completed the description of the deteriorating working environment in hospitals.

Discussion

178

179

180

181

182

183

184

185

186

187

188

189

190

191

192

193

194

195

196

197

198

199

200

201

202

Our study was designed to assess (i) the vascular surgeons' perceptions on the work life quality of their nursing staff, (ii) the impact of the nursing staff's quality of work life on their own serenity and concentration, (iii) the impact of the nursing staff's quality of work life on the safety of their surgical procedures. Our main observation is that according to 68% of vascular surgeons, their nursing staff experienced psychosocial stress and that for many of them, this significantly affected their serenity and concentration at work with frequent concerns about the surgical procedure safety. According to the surveyed vascular surgeons, the main sources of stress for the nursing staff were the high work pace (64%), demands for flexibility (56%), lack of anticipation of schedule changes (82%), and difficulties encountered with the hospital manager on these matters (44.6%). First, these results question the patient-to-nurse ratio adequacy that according to many studies, is critical for the nurses' quality of care (21-24) and work life (11). However, in the present study, the patient-to-nurse ratio was not associated with the nursing staff and vascular surgeons' quality of work life. This may be explained by the broad question on this issue in the survey with a categorical answer: ratio <8, between 8 and 10, between 11 and 13, and >13. Unfortunately, as this information is sensitive and subject to frequent changes, the exact ratio is not necessarily known by vascular surgeons. The French national hospital performance support agency (ANAP) published guidelines in 2015 to recommend <13-16 patients per nurse and the implication of certified assistant nurses for activities without technical procedures and without high patient turnover. However, data on the fluctuations in activity and staff members (e.g. training, absenteeism, burn-out), the organization of the different wards (day-to-night and week-to-weekend shift), the different activity types per nurse, the hospital budgetary situation, and the populations served are missing, although they represent critical information for the organizational equation (26,27). Many healthcare systems report that healthcare services are currently experiencing healthcare workforce shortages (14), particularly in nursing human resources (28). In our survey, some surgeons said that "although the medical and nursing

8

schools do not report a vocational crisis and have a high number of enrollments, they experience an alarming

number of dropouts during the training program because of discouragement when faced with the unpredictable nature of the working conditions or workload, particularly in conventional wards". To address this issue, the French senate recently voted in favor of a minimum patient-to-nurse ratio to improve the nurses and care assistants' working conditions (https://www.senat.fr/leg/tas22-047.html in French). By December 2024, the French healthcare authority (HAS) must determine in public hospitals, the minimum patient-to-nurse ratio per number of open beds or per number of visits (for outpatient activities) for each specialty and each activity type that will have to be implemented within 2 years.

Second, our results highlight the importance of the quality of the interactions between healthcare staff and hospital managers. Good interpersonal relationships between physicians, nurses and managers are critical for the patient safety and quality of care (29) because the medical staff's work is affected when working conditions are not good (25). In France, except for few specific and standardized situations defined by national rules, hospitals have a legal obligation to involve physicians in the review of their nursing workforce number. Yet, 66.1% of the surveyed vascular surgeons said that they had not been involved by hospital managers in these discussions. Moreover, healthcare professionals and managers often have antagonistic constraints (15) and visions of hospital activities as highlighted by one respondent who stated that "they do not have the same definition of workload". Additionally, the administrative rules for workforce allocation depend on many criteria. For instance, the rules and health economic evaluation differ when a hospital is adapting to the changes in care offers or when a hospital is planning the development of a new activity for their strategy plan. However, whatever the activity, it has been demonstrated that investments in nursing reap rewards (23,24). Small additional costs for better nursing improve patient outcomes and consequently lessen the burden and cost of post-operative care in primary care.

Third, although the vascular surgeons' serenity and concentration were somewhat affected by the nursing staff psychosocial stress, the safety of their practice did not seem to be compromised. This observation is consistent with a recent study on the vascular surgeons' capacity for resilience (31) due to their training and

their adherence to the Hippocratic oath. A more qualitative analysis with measured outcomes for patients, such as length of stay and complication rate, is needed to confirm this interpretation.

230

231

232

233

234

235

236

237

238

239

240

241

242

243

244

245

246

228

229

The main limitation of this study is its voluntary framework that could have influenced the sample representativeness. However, the French society of vascular and endovascular surgery includes almost 500 members who are representative of the French vascular surgeons in terms of age, specialty, and hospital type (public- and private-sector) where they work. The 25% response-rate represented a reasonable sample size for statistical analysis. Our sample was representative in terms of specialty, surgical practice seniority, and type of hospital. A second limitation is that respondents were volunteers, therefore strong opinions on their quality of work life may have been overestimated. However, the respondents were at first unaware that the survey would concern their nursing staff's wellbeing and this may be in favor of an accurate assessment of the nursing staff's psychosocial stress. A third limitation is that the survey did not address the seniority and stability of the nursing team and this may explain the lack of significance of the patient-to-nurse ratio. Indeed, compared with unexperienced nurses, experienced nurses should be able to manage more patients in a shorter time frame, especially in an environment with a frequent turnover. Lastly, although the present quantitative survey included open-text comments, our research would benefit from a qualitative approach with semi-structured interviews to better describe the hospital healthcare professionals' working conditions and experiences. In addition, besides the quantitative survey of vascular surgeons, a mixed-method approach that included also the nursing staff would complete and enhance our research findings.

247

248

249

250

251

Conclusions

This pilot study demonstrated that concomitantly with the ambulatory shift accelerated by the COVID-19 pandemic, the vascular surgeons' perception of their working environment is deteriorating. Workload has increased, especially in conventional surgery wards where healthcare workers encounter more patients with

many comorbidities. The work life quality for both surgeons and paramedical staff is worsening and this affects the quality of care. Working conditions and emphasis on the patient-to-nurse ratio are critical issues. When nursing staff resizing is needed, it is urgent to respect the legal frameworks that require a multidisciplinary consultation of professional experts in the field and the involvement of the concerned physicians to guarantee a safe and adequate workforce distribution. These qualitative statements complete the description of what is becoming a critical public health issue.

Acknowledgements

The authors would like to acknowledge the French society of vascular and endovascular surgery (SCVE) for the opportunity to propose the survey to their members. The authors also acknowledge the Program in Global Health at the Columbia Vagelos College of Physicians and Surgeons and Mailman School of Public Health for the coordination and funding of the research assistants. The authors thank Elisabetta Andermarcher for editing and proofreading this manuscript.

Author contributions

LD designed the study, collected the data, and reviewed and edited the manuscript; YY and SW completed the literature review, and reviewed and edited the manuscript; NLM performed the statistical analysis, interpreted the results, and wrote the original draft of the manuscript. All authors have read and agreed to the published version of the manuscript.

Competing interests

The authors have no competing interests to declare.

Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

276 References

- Melliere D, Natali J. Training of Vascular Surgeons in France: Ten Years of the French College of Vascular
 Surgery. Ann Vasc Surg. 1989 Oct;3(4):370–3.
- 2. Hingorani AP, Ascher E, Marks N, Shiferson A, Puggioni A, Tran V, et al. 219 vascular fellows' perception of the future of vascular surgery. Ann Vasc Surg. 2009;23(4):453–7.
- Le Meur N, Padilla C, Ghoroubi N, Lamirault G, Chatellier G, Gouëffic Y. Geographical Ambulatory
 Endovascular Revascularisation Disparities in France From 2015 to 2019. Eur J Vasc Endovasc Surg. 2022
 Jun;63(6):890–7.
- Jan S, Raux M, Ghoroubi N, Haulon S, Fabre D, Goueffic Y, et al. Re-hospitalization after the ambulatory
 endovascular treatment of the arteries of the lower extremities: Data of the SNDS (National System of
 Health Data) between 2013 and 2016. Ann Vasc Surg. 2022 Oct 1;68:112–3.
- 5. Hauguel A, Maurel B, Bague N, Gouaillier-Vulcain F, Costargent A, Chaillou P, et al. Management of
 ambulatory (day case) endovascular procedures for peripheral arterial disease. J Cardiovasc Surg
 (Torino). 2017 Apr;58(2):293–304.
- 6. Ali TZ, Lehman EB, Aziz F. Unplanned return to operating room after lower extremity endovascular
 intervention is an independent predictor for hospital readmission. J Vasc Surg. 2017 Jun;65(6):1735 1744.e2.
- Zeron J. Lin PH, Yang KH, Kollmeyer KR, Uceda PV, Ferrara CA, Feldtman RW, et al. Treatment outcomes and lessons learned from 5134 cases of outpatient office-based endovascular procedures in a vascular surgical practice. Vascular. 2017 Apr;25(2):115–22.
- 296 8. Oskui PM, Kloner RA, Burstein S, Zhiroff K, Kartub BR, Economides C, et al. The Safety and Efficacy of 297 Peripheral Vascular Procedures Performed in the Outpatient Setting. J Invasive Cardiol. 2015;27(5):7.
- 9. Breton M, Marshall EG, Deslauriers V, Smithman MA, Moritz LR, Buote R, et al. COVID-19 an opportunity
 to improve access to primary care through organizational innovations? A qualitative multiple case study
 in Quebec and Nova Scotia (Canada). BMC Health Serv Res. 2022 Jun 8;22(1):759.
- 10. Lee KS, Ranganathan S, Choong AMTL, Ng JJ. A scoping review on the changes in vascular surgical practice during the early phases of the COVID-19 pandemic. Semin Vasc Surg. 2021 Sep;34(3):63–73.
- 303 11. Boschetti GA, Di Gregorio S, Mena Vera JM, Pane B, Spinella G, Palombo D, et al. COVID-19 Impact on
 304 Vascular Surgery Practice: Experience From an Italian University Regional Hub Center for Vascular
 305 Pathology. Ann Vasc Surg. 2021 Jul;74:73–9.
- 12. Piazza M, Xodo A, Squizzato F, Conti GD, Boemo D, Carretta G, et al. The challenge of maintaining
 necessary vascular and endovascular services at a referral center in Northern Italy during the COVID-19
 outbreak. Vascular. 2021 Aug;29(4):477–85.
- 309 13. Birch S, O'Brien-Pallas L, Alksnis C, Tomblin Murphy G, Thomson D. Beyond demographic change in
 310 human resources planning: an extended framework and application to nursing. J Health Serv Res Policy.
 311 2003 Oct 1;8(4):225–9.
- 14. Haakenstad A, Irvine CMS, Knight M, Bintz C, Aravkin AY, Zheng P, et al. Measuring the availability of human resources for health and its relationship to universal health coverage for 204 countries and

- territories from 1990 to 2019: a systematic analysis for the Global Burden of Disease Study 2019. The
- 315 Lancet. 2022 Jun;399(10341):2129–54.
- 15. Aiken LH, Lasater KB, Sloane DM, Pogue CA, Fitzpatrick Rosenbaum KE, Muir KJ, et al. Physician and
- Nurse Well-Being and Preferred Interventions to Address Burnout in Hospital Practice: Factors
- Associated With Turnover, Outcomes, and Patient Safety. JAMA Health Forum. 2023 Jul 7;4(7):e231809.
- 319 16. Schwartz SP, Adair KC, Bae J, Rehder KJ, Shanafelt TD, Profit J, et al. Work-life balance behaviours cluster
- in work settings and relate to burnout and safety culture: a cross-sectional survey analysis. BMJ Qual Saf.
- 321 2019 Feb;28(2):142-50.
- 17. Kansoun Z, Boyer L, Hodgkinson M, Villes V, Lançon C, Fond G. Burnout in French physicians: A
- 323 systematic review and meta-analysis. J Affect Disord. 2019 Mar 1;246:132–47.
- 18. Hall LH, Johnson J, Watt I, Tsipa A, O'Connor DB. Healthcare Staff Wellbeing, Burnout, and Patient Safety:
- 325 A Systematic Review. PLOS ONE. 2016 Jul 8;11(7):e0159015.
- 326 19. Wilkinson CS, Hite KJ. Nurse-physician collaborative relationship on nurses' self-perceived job
- 327 satisfaction in ambulatory care. Lippincotts Case Manag. 2001 Mar;6(2):68–78.
- 328 20. Bontemps G. Développement de la chirurgie ambulatoire en France : plus de contraintes pour plus de
- performance ?Development of day surgery in France: More constraints for more performance? 2014
- 330 Mar;43(3):275–7.
- 21. Aiken LH, Cimiotti JP, Sloane DM, Smith HL, Flynn L, Neff DF. Effects of nurse staffing and nurse
- education on patient deaths in hospitals with different nurse work environments. Med Care. 2011
- 333 Dec;49(12):1047–53.
- 22. Rafferty AM, Clarke SP, Coles J, Ball J, James P, McKee M, et al. Outcomes of variation in hospital nurse
- staffing in English hospitals: cross-sectional analysis of survey data and discharge records. Int J Nurs Stud.
- 336 2007 Feb;44(2):175–82.
- 337 23. McHugh MD, Aiken LH, Sloane DM, Windsor C, Douglas C, Yates P. Effects of nurse-to-patient ratio
- 338 legislation on nurse staffing and patient mortality, readmissions, and length of stay: a prospective study
- in a panel of hospitals. The Lancet. 2021 May;397(10288):1905–13.
- 24. Lasater KB, McHugh M, Rosenbaum PR, Aiken LH, Smith H, Reiter JG, et al. Valuing hospital investments
- in nursing: multistate matched-cohort study of surgical patients. BMJ Qual Saf. 2021 Jan;30(1):46–55.
- 342 25. Aiken LH, Clarke SP, Sloane DM, Sochalski J, Silber JH. Hospital nurse staffing and patient mortality, nurse
- burnout, and job dissatisfaction. JAMA. 2002 Oct 23;288(16):1987–93.
- 344 26. Harper PR. A Framework for Operational Modelling of Hospital Resources. Health Care Manag Sci.
- 345 2002;5(3):165-73.
- 346 27. Sermeus W, Aiken LH, Van den Heede K, Rafferty AM, Griffiths P, Moreno-Casbas MT, et al. Nurse
- forecasting in Europe (RN4CAST): Rationale, design and methodology. BMC Nurs. 2011 Apr 18;10:6.
- 348 28. Tamata AT, Mohammadnezhad M. A systematic review study on the factors affecting shortage of nursing
- workforce in the hospitals. Nurs Open. 2023 Mar;10(3):1247–57.
- 29. Vahey DC, Aiken LH, Sloane DM, Clarke SP, Vargas D. Nurse burnout and patient satisfaction. Med Care.
- 351 2004 Feb;42(2 Suppl):II57-66.

352 353	30.	Atkinson MK, Singer SJ. Managing Organizational Constraints in Innovation Teams: A Qualitative Study Across Four Health Systems. Med Care Res Rev. 2021 Oct;78(5):521–36.
354 355 356	31.	Harkin D, Alzayyat S, Kiernan A, Ryan Á, Boland F, Renton S, et al. Vascular Surgeon Burnout and Resilience in the United Kingdom: A Report from the Vascular Society Workforce Committee. Ann Vasc Surg. 2022 Aug;84:169–78.
357		
358		
359		
360		
361		
362		
363		
364		
365		
366		
367		
368		
369		
370		
371		
372		
373		

Table I. Surgeon's perception of nursing staff's unhappiness and their work environment, in France in 2022

N	lursing staff's unha	• •	
Work environment	Sometimes N = 40*	Often N = 85*	p-value†
Hospital status			0.048
non-profit	5 (12%)	2 (2.4%)	
private	22 (55%)	44 (52%)	
public	13 (32%)	39 (46%)	
Vascular procedure			0.3
arterial only	4 (10%)	14 (16%)	
mixed	36 (90%)	71 (84%)	
Proportion of outpatient su	= -		0.5
0%	2 (5.1%)	2 (2.4%)	
<10%	18 (46%)	41 (48%)	
10-29%	9 (23%)	14 (16%)	
30-49%	2 (5.1%)	13 (15%)	
50-70%	4 (10%)	9 (11%)	
>70%	4 (10%)	6 (7.1%)	
Impact of COVID-19 on ou	tpatient activity		>0.9
decrease	2 (5.0%)	7 (8.2%)	
no impact	15 (38%)	35 (41%)	
increase	18 (45%)	33 (39%)	
do not know	5 (12%)	10 (12%)	
Impact of COVID-19 on in	patient activity		0.045
decrease	9 (22%)	38 (45%)	
no impact	26 (65%)	39 (46%)	
increase	5 (12%)	8 (9.4%)	
Adequate supervision			0.002
poor	14 (35%)	57 (67%)	
adequate	20 (50%)	22 (26%)	
do not know	6 (15%)	6 (7.1%)	
Demand for flexibility	10 (25%)	60 (71%)	<0.001
Early notice of planning ch	anges		<0.001
yes	14 (35%)	15 (18%)	
no	10 (25%)	53 (62%)	
do not know	16 (40%)	17 (20%)	
Discussion with hospital acunhappiness	dministration on nu	` '	0.058
yes	7 (18%)	13 (15%)	
sometimes	17 (42%)	22 (26%)	
no	14 (35%)	49 (58%)	
do not know	2 (5.0%)	1 (1.2%)	

^{*}n (%) - †Fisher's exact test; Pearson's Chi-squared test

Table II. Surgeon's perception of nursing staff's unhappiness by surgery setting, in France in 2022.

		Outpatient			Inpatient	
	Sometimes, N = 33*	Often N = 85*	p-value†	Sometimes N = 33*	Often N = 85*	p-value†
Change in workloa	d since ambula	atory shift	0.6			0.073
decrease	2 (6.1%)	5 (5.9%)		9 (27%)	9 (11%)	
no change	3 (9.1%)	16 (19%)		8 (24%)	19 (22%)	
increase	20 (61%)	44 (52%)		8 (24%)	38 (45%)	
do not know	8 (24%)	20 (24%)		8 (24%)	19 (22%)	
Patient-to-nurse ra	tio		>0.9			0.2
<8	5 (21%)	17 (27%)		6 (19%)	6 (7.2%)	
8-13	13 (56%)	34 (56%)		19 (59%)	61 (73.5%)	
>13	5 (22%)	10 (16%)		7 (22%)	16 (19.3%)	
Number of specials	ties per nurse		0.8			0.2
1	1 (3.1%)	2 (2.4%)		11 (33%)	13 (15%)	
2-3	3 (9.4%)	6 (7.1%)		13 (39%)	44 (52%)	
4-5	11 (34%)	36 (43%)		6 (18%)	21 (25%)	
>5	17 (53%)	40 (48%)		3 (9.1%)	7 (8.2%)	
Work pace			0.006			<0.001
low to adequate	21 (64%)	27 (32%)		19 (58%)	15 (18%)	
sustained	7 (21%)	38 (45%)		13 (39%)	65 (76%)	
do not know	5 (15%)	20 (24%)		1 (3.0%)	5 (5.9%)	
Lack of nursing ca	re due to time o	constraints	0.7			0.2
0-15%	17 (52%)	33 (39%)		13 (39%)	30 (35%)	
16-30%	5 (15%)	13 (15%)		10 (30%)	22 (26%)	
>30%	2 (6.1%)	8 (9.4%)		1 (3.0%)	15 (18%)	
do not know	9 (27%)	31 (36%)		9 (27%)	18 (21%)	

^{*}n (%)

[†]Fisher's exact test; Pearson's Chi-squared test

Table III. Factors associated with nursing staff's unhappiness, in France in 2022. Method: multivariate

logistic regression with often vs. sometimes unhappy

Characteristics	OR*	95% CI†	p-value
Impact of COVID-19 on conventional activity			
no impact	_	_	
decrease	2.99	0.97, 10.2	0.065
increase	0.46	0.08, 2.66	0.4
Demand for flexibility			
no	_	_	
yes	5.03	1.74, 15.8	0.004
Work pace in conventional wards			
low to adequate pace	_	_	
sustained pace	4.71	1.66, 14.1	0.004
do not know	27	2.29, 801	0.02
Early notice of planning changes			
yes	_	_	
no	4.50	1.27, 16.9	0.021
do not know	1.23	0.33, 4.62	0.8

^{*}OR = Odds Ratio

[†]CI = Confidence Interval

Table IV. Factors associated with vascular surgeons' perturbed work safety in conventional wards, in

France in 2022. Method: univariate analysis.

Characteristics	Rarely N = 21*	Occasionally N = 61*	Very frequently N = 33*	p-value†
Patient-to-nurse ratio				0.6
<8	2 (9.5%)	8 (13%)	2 (6.1%)	
8-13	15 (71%)	44 (72%)	22 (67%)	
>13	4 (19%)	9 (15%)	9 (27%)	
Number of specialties per nurse				0.9
1	5 (24%)	9 (15%)	6 (18%)	
2-3	11 (52%)	33 (54%)	14 (42%)	
4-5	4 (19%)	13 (21%)	9 (27%)	
>5	1 (4.8%)	6 (9.8%)	4 (12%)	
Work pace				0.015
low to adequate pace	11 (52%)	17 (28%)	4 (12%)	
sustained pace	10 (48%)	42 (69%)	26 (79%)	
do not know	0 (0%)	2 (3.3%)	3 (9.1%)	
Lack of nursing care due to time constraints				0.12
0-15%	11 (52%)	23 (38%)	9 (27%)	
16-30%	4 (19%)	22 (36%)	7 (21%)	
>30%	1 (4.8%)	7 (11%)	8 (24%)	
do not know	5 (24%)	9 (15%)	9 (27%)	
Inpatient increased comorbidities	8 (38%)	42 (69%)	23 (70%)	0.028
Nursing staff's unhappiness				<0.001
sometimes	12 (57%)	21 (34%)	2 (6.1%)	
often	9 (43%)	40 (66%)	31 (94%)	

^{*}n (%)

[†]Fisher's exact test; Pearson's Chi-squared test

Table V. Factors associated with vascular surgeons' perturbed work safety in conventional wards, in France

in 2022. Method: Multivariate multinomial analysis - Reference: Safety of care rarely perturbed.

		Occasionally	1	•	Very frequent	ly
Characteristics	OR*	95% CI†	p-value	OR*	95% CI†	p-value
Patient-to-nurse ratio						
<8	_			_	_	
8-13	2.5	0.38, 16.4	0.3	2.73	0.24, 30.6	0.4
>13	0.82	0.19, 3.50	8.0	1.88	0.38, 9.28	0.4
Work pace						
low to adequate pace		_		_	_	
sustained pace	1.82	0.55, 5.95	0.3	2.61	0.54, 12.5	0.2
Inpatient increased comorbidities						
no	_	_		_	_	
yes	3.92	1.27, 12.1	0.018	4.75	1.21, 18.6	0.026
Nursing staff's unhappiness						
sometimes	_	_		_	_	
often	2.5	0.77, 8.18	0.13	16.3	2.58, 102	0.003

^{*}OR = Odds Ratio

[†]CI = Confidence Interval

wards, in France in 2022. Method: univariate analysis.

Characteristics	Rarely N = 16*	Occasionally N = 49*	Very frequently N = 43*	p-value†
Inpatient increased comorbidities	7 (44%)	29 (59%)	33 (77%)	0.042
Patient-to-nurse ratio				0.1
<8	14 (88%)	33 (67%)	28 (65%)	
8-13	1 (6.2%)	8 (16%)	2 (4.7%)	
>13	1 (6.2%)	8 (16%)	13 (30%)	
Number of specialties per nurse				0.7
1	4 (25%)	10 (20%)	6 (14%)	
2-3	6 (38%)	26 (53%)	19 (44%)	
4-5	5 (31%)	9 (18%)	12 (28%)	
>5	1 (6.2%)	4 (8.2%)	6 (14%)	
Work pace				0.092
low to adequate pace	7 (44%)	17 (35%)	8 (19%)	
sustained pace	9 (56%)	32 (65%)	35 (81%)	
Lack of nursing care due to time constra	nints			0.028
0-15%	7 (44%)	22 (45%)	12 (28%)	
16-30%	1 (6.2%)	15 (31%)	14 (33%)	
>30%	1 (6.2%)	4 (8.2%)	10 (23%)	
do not know	7 (44%)	8 (16%)	7 (16%)	
Nursing staff's unhappiness				0.006
sometimes	7 (44%)	21 (43%)	6 (14%)	
often	9 (56%)	28 (57%)	37 (86%)	
Discussion with hospital administration				0.002
on nursing staff's unhappiness	0 (400()	04 (400()	07 (000()	3.002
no	2 (12%)	24 (49%)	27 (63%)	
yes	8 (50%)	6 (12%)	4 (9.3%)	
sometimes	6 (38%)	18 (37%)	10 (23%)	
to not know	0 (0%)	1 (2.0%)	2 (4.7%)	

^{*}n (%)

[†]Fisher's exact test; Pearson's Chi-squared test

Table VII. Factors associated with vascular surgeons' perturbed serenity and concentration in conventional wards, in France in 2022. – Method: Multivariate multinomial logistic regression - Reference: serenity and concentration during care is rarely impaired -.

		Occasiona	lly		Very freque	ently
Characteristics	OR*	95% CI†	p-value	OR*	95% CI†	p-value
Discussion with hospital administration on nursing staff's unhappiness						
yes, or sometimes	_			_	_	
no	6.78	1.33, 34.6	0.022	10.6	1.97, 57.0	0.006
Inpatient increased comorbidities						
no	_			_	_	
yes	1.42	0.43, 4.77	0.6	4.64	1.17, 18.5	0.03
Nursing staff's unhappiness						
sometimes	_			_	_	
often	0.86	0.25, 2.94	8.0	4.78	1.06, 21.6	0.042

^{*}OR = Odds Ratio,

[†]CI = Confidence Interval

432	Supplementary material for Destrieux et al. Work environment differences between
433	outpatient and inpatient surgery: a pilot study on the vascular surgeons' perceptions.
434	
435	Survey on the quality of work life for vascular surgeons
436	
437	I – Hospital characteristics
438	
439 440 441 442 443 444 445	 1- What is the legal status of your hospital? □ Public-sector university hospital □ Private-sector hospital □ Non-profit sector hospital □ Other
446 447 448 449 450 451	2- What vascular surgery procedures do you perform? ☐ Arterial and venous ☐ Arterial exclusively ☐ Venous exclusively ☐ Other
452	3- Today, what proportion of arterial surgery procedures do your carry out in the day surgery ward?
453 454 455 456 457 458 459	□ >90% □ >70% □ 50-70% □ 30-49% □ 10-29% □ <10%
460 461	4- How many different medical-surgical specialties does a nurse in a conventional surgery ward have to deal with at the same time?
462 463 464 465 466	□ 1 □ 2 to 3 □ 4 to 5 □ >5
467 468	5- How many different medical-surgical specialties does a nurse in a day surgery ward have to deal with at the same time?
469 470 471	□ 1 □ 2 to 3 □ 4 to 5

473	6- What is the current patient-to-nurse ratio in your conventional surgery ward?
474 475 476 477 478 479 480	 <8 8 to 10 11 to 13 >13 Do not know. Other 7- What is the current patient-to-nurse ratio in your day surgery ward?
481 482 483 484 485 486 487	 □ <8 □ 8 to 10 □ 11 to 13 □ >13 □ Do not know. □ Other
488	II – Impact of the COVID-19 crisis on the vascular surgery activity changes
489	
490	8- What impact did the COVID-19 crisis have on the activities of your vascular surgery wards?
491 492 493 494 495 496 497	 □ Increase of outpatient surgery procedures AND no impact on inpatient surgery procedures. □ No impact. □ Increase of both outpatient AND inpatient surgery procedures. □ Increase of outpatient surgery procedures AND decrease of inpatient surgery procedures. □ No impact on outpatient surgery procedures AND decrease of inpatient surgery procedures. □ Do not know.
498 499 500	9- In your opinion, what is the impact of the increase of outpatient surgical procedure on the disease burder (comorbidities, complexity of surgical management) of patients hospitalized in conventional and day surgery wards?
501 502 503 504 505 506	 □ No impact on inpatient and outpatient burden □ Increased burden of both inpatients and outpatients □ Increased burden of inpatients only □ Increased burden of outpatients only □ Do not know.
507 508	10 – In the context of a recent opening of a day vascular surgery ward, what is the impact on the delivery of follow-up and treatment information? (Multiple answers possible)
509 510 511 512 513 514 515	 □ Development of new materials to inform patients. □ Materials to inform patients are not optimal. □ Already optimal materials to inform patients. □ More compliant patients in day surgery wards than in conventional wards □ Less compliant patients in day surgery wards than in conventional wards □ No impact □ Too soon to observe any impact

517	III - Changes in surgical activities and adequacy of nursing resources (nurses and certified assistant nurses)
518	
519 520	11- In your opinion, what impact did the opening or increase of outpatient surgery activity have on the nursing staff's workload in your vascular surgery wards?
521 522 523 524 525 526 527 528 529 530 531 532 533 534 535	 □ No impact □ Workload increase in both day AND conventional vascular surgery wards. □ Workload increase in the day vascular surgery ward AND decrease in the conventional vascular surgery ward. □ Workload increase in the day vascular surgery ward AND no impact in the conventional vascular surgery ward. □ Workload decrease in both day AND conventional vascular surgery wards. □ Workload decrease in the day vascular surgery ward AND no impact in the conventional vascular surgery ward. □ Workload decrease in the day vascular surgery ward AND increase in the conventional vascular surgery ward. □ No impact in the day vascular surgery ward AND increase in the conventional vascular surgery ward. □ No impact in the day vascular surgery ward AND decrease in the conventional vascular surgery ward. □ Do not know.
536 537	12- In your opinion, is the nursing staff' work pace currently adapted to manage your outpatients (including technical care, listening time and information time)?
538 539 540 541 542	☐ Slow ☐ Adapted ☐ Fast ☐ Do not know.
543 544	13- In your opinion, is the nursing staff's work pace currently adapted to manage your inpatients (including technical care, listening time and information time)?
545 546 547 548 549	☐ Slow ☐ Adapted ☐ Fast ☐ Do not know.
550 551 552	14- In your opinion, what percentage of your inpatients was affected at least once by lack of nursing care due to the nurses' lack of time in 2022 (e.g. technical procedure not performed, prescribed treatment not administered, information not provided)?
553 554 555 556 557 558 559 560 561	□ 0-15% □ 16-30% □ 31-45% □ 46-60% □ 61-75% □ 76-90% □ >90% □ No impact

562 563 564	15- In your opinion, what percentage of your outpatients was affected at least once by lack of nursing card due to nurses' lack of time in 2022 (e.g. technical procedure not performed, prescribed treatment not administered, information not provided)?
565 566 567 568 569 570 571 572 573	□ 0-15% □ 16-30% □ 31-45% □ 46-60% □ 61-75% □ 76-90% □ >90% □ No impact
574	IV - Nursing staff's quality of work life
575	16 - In your opinion, are nursing staff properly supervised in your vascular surgery wards?
576 577 578 579	□ No □ Yes □ Do not know
580 581	17- In your opinion, do nursing staff have to change tasks, positions, or functions at a moment's notice to meet current constraints?
582 583 584 585 586	 No, rarely Yes, sometimes Yes, often Do not know
587 588	18 - In your opinion, are nursing staff informed sufficiently in advance about their working hours or about changes to their work schedule?
589 590 591 592	 □ No, rarely □ Yes, most of the time □ Do not know
593 594	19- In your opinion, are nursing staff unhappy with their working conditions (high levels of occupational stress, anxiety, unsatisfaction)?
595 596 597 598	□ No, rarely□ Yes, most of the time□ Do not know
599	
600	
601	
602	

603	20- In your opinion, this unhappiness is most often related to:
604 605 606 607 608 609 610 611 612 613	 □ Work overload □ Lack of professional recognition □ Physicians' unsuitable or oppressive attitude □ Hospital managers' unsuitable or oppressive attitude □ Poor care management organization □ Relationship problems with other healthcare staff members □ Inadequate pay □ I do not perceive unhappiness in my team □ I do not know
614	V - Impact of the nursing staff's quality of work life on the vascular surgeons' working conditions
615	
616 617	21- As a surgeon, do you have the means to discuss with your administration in the event of unhappiness or difficulties among nursing staff?
618 619 620 621 622	☐ No ☐ Sometimes ☐ Yes ☐ I do not know
623 624	22- How would you describe your healthcare human resources currently working in the conventional and dasurgery wards? Multiple answers possible
625 626 627 628 629 630 631	 □ Very stable: little turnover □ Long-term bed closures due to unreplaced departures □ Frequent use of temporary staff to keep beds open □ Constant turnover due to regular departures that are easily replaced without temporary staff □ Fragilized by frequent unreplaced departures. □ Other
632 633	23- Are your serenity and professional concentration affected by difficulties related to nursing staff (organization, skills, experience)?
634 635 636 637 638 639	 □ Rarely □ Occasionally □ Very frequently □ Do not know □ Other
640	24- Is the safety of your surgical procedures affected by problems with nursing staff?
641 642 643 644 645 646	 □ Rarely □ Occasionally □ Very frequently □ Do not know □ Other

647	VI - Opportunities given to surgeons to get involved in the hospital care organization.
648 649	25- Can you or your medical representatives effectively express your opinions and proposals to the hospita management team when the staff organization or workload appears to be detrimental to care?
650 651 652 653 654 655 656 657	 □ Never □ With difficulty □ Occasionally □ Often □ Always □ Do not know □ Other
658 659	26- Have you or your medical representatives been asked by your hospital management team to participate in the review of healthcare staffing requirements in your day surgery and conventional surgery wards?
660 661 662 663 664 665 666	 Yes, optimal staffing levels are reviewed yearly with the management committee. Yes, optimal staffing levels are part of every hospital project renewal. Yes, this is done when there is a significant change in surgical activity or a new type of activity. No, to my knowledge, the healthcare staffing review does not involve the physicians in my hospital Do not know Other
667 668	27- In your opinion, are changes in the medical care organization anticipated, discussed, and clearly explained to practitioners by the hospital management team?
669 670 671 672 673 674	 □ Never □ Occasionally □ Often □ Always □ Do not know
675	28- What is the quality of your relations with your hospital management team?
676 677 678 679 680 681	 □ Not existent □ Poor □ Unstable □ Satisfactory □ Excellent □ Other
682	